

Construction of a *Populus tremuloides* Michx. BAC library

By M. FLADUNG¹⁾, H. KAUFMANN²⁾, T. MARKUSSEN and H. HOENICKA

Federal Research Centre for Forestry and Forest Products, Institute for Forest Genetics and Forest Tree Breeding, Sieker Landstr. 2, D-22927 Grosshansdorf, Germany

(Received 13th November 2006)

Abstract

We have constructed an aspen (*Populus tremuloides* Michx., line Turesson141) BAC library containing 55,296 clones in total. A random sampling of 86 BAC clones indicated an average insert size of 76 kb with a range of 20 to 160 kb. Twelve percent of the BAC clones in the library have an insert size larger than 100 kb. Based on an estimated genome size for *Populus* of 500 Mbp, library coverage is about 8 haploid genome equivalents. This library will be screened using AFLP marker identified before co-segregating with gender in a *P. tremula* x *P. tremuloides* progeny, where Turesson141 was the male parent.

Key words: Bacterial Artificial Chromosome, *Populus*, genome size, aspen.

Introduction

Poplar is one of the world's most important tree species. Their rapid growth and capacity to supply a broad range of wood products has led to their widespread cultivation in Europe and North America (DICKMAN and STUART, 1983; STETTLER et al., 1996). In addition to their commercial value, poplars and hybrid poplars are ecologically important as they are effective in the control of nutrients and toxic compounds found in agricultural liquid waste and landfill effluents (BRAATNE, 1999).

The genus *Populus* serves as model tree species for gene technological studies and genomic research for several reasons: (1) *Populus trichocarpa* is the only tree species which has been completely sequenced up to now (<http://www.ornl.gov/sci/ipgc/>), (2) The genome size of poplar with approximately 500 Mbp is small. This is similar in size to the rice genome and only four times larger than the genome of *Arabidopsis*, yet 40 times smaller than the genome of pines (3). Further, there are many genomic efforts in this genus including availability of a high number of expressed sequence tag (EST-) data (BHALLERAO et al., 2003).

However, as for most of the forest tree species improvement in poplar is hampered by tree-specific features like long juvenility phases and slow growth. Despite of successful sequencing in one poplar species,

very few knowledge is available about tree genetics due to the lack of inbred lines required for intensive genetic studies. Nowadays, novel molecular breeding tools have been developed allowing the creation of and screening for novel poplar genotypes.

During the last decade several genomic maps have been published for different poplar species using RAPD-, AFLP-, EST-, and SSR marker (for overview: CERVERA et al., 2004, e.g. *P. trichocarpa* (CERVERA et al., 2001), *P. alba* (YIN et al., 2001), *P. nigra* (CERVERA et al., 2001), *P. deltoides* (YIN et al., 2002), and *P. tremula* x *P. tremuloides*, *P. tremula* x *tremula* (MEYER et al., pers. communication). On basis of these maps a physical mapping and cloning of genes associated with important developmental and/or wood traits would be feasible if a BAC-(Bacterial Artificial Chromosome) library is available.

For woody plants the construction of BAC libraries was reported mainly for fruit trees and horticultural crops, like coffee (NOIR et al., 2004), apple (VINATZER et al., 1998; PATOCCHI et al., 1999; XU et al., 2001), peach (WANG et al., 2001), apricot (VILANOVA et al., 2003), grape (TOMKINS et al., 2001) and rose (KAUFMANN et al., 2003). But also for forest trees BAC libraries have been constructed for Eucalyptus (GRATTAPAGLIA, 2004) and *Liriodendron tulipifera* (<http://www.greenbac.org/index.html>). Two BAC libraries have been constructed for *P. trichocarpa* in order to characterise and isolate the MXC3 locus a major gene for resistance to poplar leaf rust (STIRLING et al., 2001) as well as the R-locus for resistance against *Melampsora larici-populina* causing leaf rust (FAIVRE RAMPANT et al., 2006). The BAC library from the *P. trichocarpa* clone 383-2499 (Nisqually-1) was used to sequence the whole poplar genome. In frame of the EU-Network of Excellence "EvolTree" (www.evoltree.org) it is planned to construct a BAC library from *Quercus* but also further ones from *Populus*.

In this study we report the construction and characterisation of a BAC library from aspen-*Populus tremuloides*. To the best of our knowledge, the *P. tremuloides* (aspen) BAC library reported here is the first large-insert DNA library so far constructed for this species, and the third reported for the genus *Populus*.

Material and Methods

Plant material

Leaves of a single, approximately 40 years-old *Populus tremuloides* Michx. tree (clone Turesson141) growing in the 'Arboretum Tannenhöft' of the Institute of Forest Genetics and Forest Tree Breeding, Grosshansdorf, Ger-

¹⁾ Corresponding author: Matthias Fladung, Federal Research Centre for Forestry and Forest Products, Institute for Forest Genetics and Forest Tree Breeding, Sieker Landstr. 2, D-22927 Grosshansdorf. Fax: +49-4102-696200, Email: mfladung@uni-hamburg.de.

²⁾ Present address: Leibniz-University, Institute of Floriculture and Tree Nursery Science, Herrenhäuser Str. 2, D-30419 Hanover, Germany.

many, were used. Leaf material of different ages (young [starting to unfold] and mature [up to four weeks old]) was harvested, kept for two days in the dark, frozen in liquid nitrogen and stored at -80°C . For the isolation of nuclei suitable for BAC cloning frozen leaf material was ground under liquid nitrogen using the Retsch Bead Mill (Retsch, Duesseldorf, Germany) or mortar and pestle.

BAC library construction

The aspen BAC library was constructed following the protocol described by KAUFMANN et al. (2003). However, each working step had to be optimised for *P. tremuloides* as described in the results section. All other molecular procedures not cited separately were done as described in SAMBROCK et al. (1989).

The isolation and lysis of aspen nuclei as well as the PMSF (Phenylmethylsulphonylfluoride) treatment was performed as described in KAUFMANN et al. (2003) with some modifications (see results). For partial restriction *Hind*III concentrations between 0.8 and 50U per agarose plug were tested to obtain DNA fragments in a size range of 100–350kb. Restricted DNA was separated using the pulsed-field gel electrophoresis CHEF-DRII-system (Bio-Rad, Germany) and DNA of the appropriate size was cut out of the gel and electroeluted (STRONG et al., 1997).

Eluted DNA was dialysed and ligated into the BAC vector pBeloBACII (KIM et al., 1996). The plasmid has a size of 7.4 kb and carries a chloramphenicol-resistance gene. Sequence of the vector as well as restriction map are available under <http://www.sanger.ac.uk/Teams/Team53/psub/sequences/pbelobac.shtml> and <http://www.sanger.ac.uk/Teams/Team53/psub/sequences/pbelobac.pdf>, respectively. The plasmid was restricted with *Hind*III, dephosphorylated and tested for cloning ability as described by PETERSON et al. (2000).

Following heat inactivation and dialysis the ligation mixtures were transformed in electrocompetent *E. coli* DH10B cells (Life technologies) using an *E. coli* Pulser transformation apparatus (BioRad, Germany).

Characterisation of the BAC library

The ligation mixtures showing a satisfactory transformation rate and a high proportion of white colonies were tested more in detail.

BAC-DNA was prepared by a standard alkaline lysis method from 2-ml overnight cultures in TB with antibiotic and resuspended in 30 μl of TE. Five μl of BAC-Mini-DNA were digested with *Not*I in a volume of 20 μl at 37°C for 2 hrs. For pulsed-field gel electrophoresis, samples were loaded on a 1% TAE agarose gel and electrophorised with 5 to 15 sec pulse time for 20 h with field strength of 4.5V/cm at 14°C .

Results and Discussion

Optimisation of nucleus preparation, restriction, and pulsed-field gel electrophoresis

The protocol for construction of a rose BAC library (KAUFMANN et al., 2003) was modified to optimize the

construction of an aspen BAC library. Rose and aspen have a similar genome size which is about four times larger than *Arabidopsis* (about 500 Mbp or a 1C value of about 0.5 pg (YOKOYA et al., 2000), <http://www.ornl.gov/sci/ipgc/>). In both genera DNA isolation might be hampered by presence of polyphenolic compounds or polysaccharides in leaves.

For nuclei and subsequent DNA isolation, in total, nine different preparations were done. Out of these, five preparations yielded in high-quality DNA which could be used for cloning experiments. Comparative grinding under liquid nitrogen using the Retsch Bead Mill and traditional mortar and pestle revealed no differences in nuclei yield, but later on a higher cloning success was obtained when using mortar and pestle.

To identify leaf age for optimal nuclei preparation young (starting to unfold) and mature (up to four weeks old) leaves were harvested. Slight or no differences were found with respect to DNA quality and purity. On the other hand, composition of lysis buffer is crucial for nuclei quality. The lysis buffer according to PETERSON et al. (2000) was slightly modified: concentrations of PVP40 and 2-mercaptoethanol were increased to 5% and 0.5%, respectively, to avoid oxidative processes which affect the quality of DNA. Increase of the Triton X 100 concentration to 1% resulted in a more effective lysis of chloroplasts, thus, a higher purity of the nuclei was achieved.

Optimal *Hind*III concentration for partial restriction to obtain DNA fragments with sizes between 100 and 350 kb was found between 4 to 40 U per agarose plug with an incubation time of 30 minutes at 37°C . Three different pulsed-field gel electrophoresis conditions were tested. First, as described for construction of the rose BAC library (KAUFMANN et al., 2003), a two-phase electrophoresis with 60 to 90 sec pulse time for 5 hours and subsequent 4 sec for 16 hours was tested. By using these electrophoresis conditions BAC clones with sizes about 50 kb were obtained. A three-phase electrophoresis with 60 to 90 sec pulse time for 8 hours, 4 sec for 20 hours, and finally 4.5 sec for 16 hours (ALLOUIS et al., 2001) yielded clones with sizes between 50 and 70 kb. Third, a one-phase electrophoresis with 1 to 40 sec pulse time for 24 hours revealed inserts between 80 and 100 kb.

A further significant point is loading density of the pulsed-field gel. It is important to obtain a minimal density of 0.5 $\mu\text{g}/\mu\text{l}$ of DNA in the eluate but without overloading the gel, the latter leading to co-migration of small DNA fragments. To slightly increase loading density the thickness of the loading slot was increased from the standard 1.5 to 4.5 mm. The eluate used in the successful cloning experiment had a concentration of 3 $\text{ng}/\mu\text{l}$.

Different conditions for ligation and transformations were tested to gain optimal number of clones and largest insertions (Table 1). For ligation amount of DNA and the vector, but as well also different ligation buffers (Invitrogen, Germany, and NBI Fermentas, Lithuania) were tested. The optimal conditions for ligation revealed to be 15 ng vectors per 100 μl in combination with as high as possible concentration of DNA in PEG-free

buffer. In the subsequent desalting procedure dialysis against 0.5 x TE (KAUFMANN et al., 2003) was found to be more effective than 10% PEG 8000 (PETERSON et al., 2000). Finally, addition of tRNA as carrier (ZHU et al., 1999) increased the transformation efficiency.

Protocol for the construction of an aspen BAC library

Four grams of frozen leaf material were grinded under liquid nitrogen and resuspended in 40 ml lysis buffer (PETERSON et al., 2000) supplemented with 5% PVP40, 1% Triton X100 as well as 0,5% 2-mercaptoethanol. Following filtration through Miracloth (Calbiochem, San Diego, USA) and various washing steps using lysis buffer the nuclei were transferred to 1 ml lysis buffer without 2-mercaptoethanol mixed with 1ml of 1.2% Low Melting Point agarose (Seaplaque FMC, USA) and moulded to small plugs.

The plugs were incubated in 0,5M EDTA pH 9.0, 1% sodiumlaurylsarcosine, 2% PVP 40 and 0.4mg/ml proteinase K for 72 hours at 50°C with two buffer changes. Following lysis, the blocks were washed five times for 30 min at room temperature in 10 mM Tris-HCl pH 7.5, 10 mM EDTA, 2% PVP 40 and 0.5 mM PMSF. Before restriction the plugs were chopped into small pieces and equilibrated for 90 min in React 2-buffer plus 4 mM spermidine-HCl and 100 µg/ml BSA. Following 15 min incubation in React 2-buffer (Invitrogen, Germany),

4 mM spermidine-HCl, 100 µg/ml BSA and 40 U *Hind*III on ice restriction was performed at 37°C for 30 min. Restriction was stopped by adding 50 mM EDTA.

Partially restricted DNA was separated through pulsed-field gel electrophoresis (CHEF-DRII-System) on a 1% preparative LMP agarose gel in 1xTAE using the following parameter: 1–40 sec pulse time, 3,5V/cm, 1200, 10°C for 20 hours. Slices of agarose containing DNA of sizes from 80 to 180 kb, from 180 to 250 kb and from 250 to 350 kb were excised and electro-eluted following the protocol by STRONG et al. (1997). Ligation with *Hind*III digested pBeloBACII-vector was performed in a final volume of 170 µl with 7.5 U T4-DNA-Ligase (Life Technologies), 450 ng eluted DNA (fraction 250 to 350 kb) and 25 ng vector. Following heat inactivation of ligase and desalting 2 µl of ligation solution were mixed with 40 µl competent *E. coli* DH10B cells (Invitrogen, Germany) using an *E. coli* pulser (Biorad, Germany) at 11 kV/cm (KAUFMANN et al., 2003). After 45 min of recovery at 37°C in 0.5 ml SOC-medium transformed *E. coli* cells were plated on LB-agar containing IPTG, X-gal (SAMBROOK et al., 1989) and 12.5 µg/ml chloramphenicol and incubated at 37°C for 20 hours. Transformation rate was about 800 recombinant (white) clones per µl ligation solution. In total 55,296 clones were picked and stored in one hundred forty-four 384-well microtiter plates.

Table 1. – Influence of different parameters on ligation and transformation.

	Efficiency of transformation
Standard ligation, volume=100µl (Kaufmann et al. 2003)	100%
50ng DNA, 20 ng vector, 5xligase buffer without PEG, 5U ligase	
50ng DNA, 20 ng vector, 5xligase buffer <u>with</u> PEG, 5U ligase	18%
<u>DNA-dilution 1:1 (25 ng DNA/100µl)</u> , 20 ng vector, 5xligase buffer without PEG, 5U ligase	46%
<u>DNA-dilution 2:1 (37 ng DNA/100µl)</u> , 20 ng vector, 5xligase buffer without PEG, 5U ligase	73%
50ng DNA, <u>30 ng vector</u> , 5xligase buffer without PEG, 5U ligase	70%
50ng DNA, <u>15 ng vector</u> , 5xligase buffer without PEG, 5U ligase	200%
Standard transformation (Petersen et al. 2000)	100%
11kV/cm, pre-treatment: dialysis 10% PEG, no carrier	
11kV/cm, pre-treatment: <u>dialysis 0,5xTE</u> , no carrier	137%
11kV/cm, pre-treatment: dialysis 0,5xTE, <u>addition of 1µg tRNA as carrier</u>	200%

* modifications underlined.




Figure 1. – Pulse-field gel electrophoresis of *NotI* digested aspen BAC clones. Inverted image of an ethidium bromide stained gel showing the inserts of 38 clones.




Figure 2. – Insert size distribution of clones in the aspen BAC library. 86 BAC clones were randomly selected and digested with *NotI*. The fragment sizes were determined by pulse-field gel electrophoresis and plotted against frequency of occurrence.

Characterisation of the aspen BAC library

To determine the insert-size distribution and average insert size of the aspen BAC library, DNA was isolated from 90 randomly selected clones, digested with *NotI* and separated by pulse-field gel electrophoresis (Fig. 1). Four clones had no insert (corresponding to 4.4%), while the residual contained inserts ranging from 20 to 160 kb in size (Fig. 2). Average size is 76 kb, and twelve percent of the BAC clones in the library have an insert size higher than 100 kb.

The percentage of clones in the library that contained organelle DNA was determined by less than 5%. Most of these clones had chloroplast sequences, but only less than 0.1% carries inserts of mitochondrial origin. Based on the haploid genome size of about 0.5 pg correspond-

ing to 500 Mbp (<http://www.ornl.gov/sci/ipgc/>), this library is estimated to contain 8 genome equivalents. Therefore, the probability of recovering any given sequence of the aspen genomic DNA from this library is higher than 99% (CLARKE and CARBON, 1976).

The library reported here is well suited for many applications in forest tree genome research. In combination with the *P. tremuloides* genomic map, a map-based cloning approach can be initiated to investigate molecular markers for gender which has already been mapped (Markussen et al., *Silvae Genetica*, in press). Considering the high number of genomic maps available for this genus as well as the genome sequence of *P. trichocarpa* (<http://www.ornl.gov/sci/ipgc/>) which has been generated from a BAC library a comparative physical mapping approach is now feasible in the genus *Populus*.

Acknowledgements

This study was supported by the German Research Foundation (FL263/2-1 and FL263/2-2). We wish to thank O. NOWITZKI and A. TUSCH for excellent technical assistance.

References

- ALLOUIS, S., X. QI, M. D. GALE and K. M. DEVOS (2001): Construction of a BAC library of pearl millet, *Pennisetum glaucum*. *Theor. Appl. Genet.* **102**: 1200–1205.
- BHALERAO, R., O. NILSSON and G. SANDBERG (2003): Out of the woods: forest biotechnology enters the genomic era. *Curr. Opin. Biotechnol.* **14**: 206–213.
- BRAATNE, J. H. (1999): Biological aspects of hybrid poplar cultivation on floodplains in western North America. Prepared for U.S. EPA (Region 10), Seattle, WA, Document No. 910-R-99-002.
- CERVERA, M. T., M. M. SEWELL, P. FAIVRE-RAMPANT, V. STORME and W. BOERJAN (2004): Genome mapping in *Populus*. In: KUMAR, S., FLADUNG, M. (Eds) *Molecular Genetics and Breeding of Forest Trees*. The Haworth Press, Binghamton, 387–410.
- CERVERA, M. T., V. STORME, B. IVENS, J. GUSMÃO, B. H. LIU, V. HOSTYN, J. VAN SLYCKEN, M. VAN MONTAGU and W. BOERJAN (2001): Dense genetic linkage maps of three *Populus* species (*Populus deltoides*, *P. nigra* and *P. trichocarpa*) based on AFLP and microsatellite markers. *Genetics* **158**: 787–809.
- CLARKE, L. and J. CARBON (1976): A colony bank containing synthetic Col EI hybrid plasmids representative of the entire *E. coli* genome. *Cell* **9**: 91–99.
- DICKMAN, D. I. and K. W. STUART (1983): *The Culture of Poplars in Eastern North America*. Department of Forestry Publications, Michigan State University, East Lansing, Michigan.
- FAIVRE RAMPANT, P., L. BATAILLE, A. BRESSON, N. BOUDET, B. CHALHOUB, V. JORGE, A. DOWKIW, V. GUERIN, J. P. MASLE, M. VILLAR and C. BASTIEN (2006): Genetic and Physical mapping of a genetic factor involving in partial rust resistance in *Populus trichocarpa*. IV International Poplar Symposium, Nanjing, China.
- GRATTAPAGLIA, D. (2004): Integrating genomics into *Eucalyptus* breeding. *Genet. Mol. Res.* **3**: 369–379.
- KAUFMANN, H., L. MATTIESCH, H. LÖRZ and T. DEBENER (2003): Construction of a BAC library of *Rosa rugosa* Thunb. And assembly of a contig spanning *Rdr1*, a gene that confers resistance to blackspot. *Mol. Genet. Genomics* **268**: 666–674.
- KIM, U. J., B. W. BIRREN, T. SLEPAK, V. MANCINO, D. BOYSEN, H. L. KANG, M. I. SIMON and H. SHIZUYA (1996): Construction and characterization of a human bacterial artificial chromosome library. *Genomics* **34**: 213–218.
- NOIR, S., S. PATHEYRON, M. C. COMBES, P. LASHERMES and B. CHALHOUB (2004): Construction and characterisation of a BAC library for genome analysis of the allotetraploid coffee species (*Coffea arabica* L.). *Theor. Appl. Genet.* **109**: 225–230.
- PATOCCHI, A., B. A. VINATZER, L. GIANFRANCESCO, S. TARTARINI, H. B. ZHANG, S. SANSVINI and C. GESSLER (1999): Construction of a 550-kb BAC contig spanning the genomic region containing the apple scab resistance gene Vf. *Mol. Genet. Genomics* **262**: 884–891.
- PETERSON, D. G., J. P. TOMKINS, D. A. FRISCH, R. A. WING and A. H. PATERSON (2000): Construction of plant bacterial artificial chromosome (BAC) libraries: an illustrated guide. *JAG 5*: www.ncgr.org/research/jag.
- SAMBROOK, J., E. F. FRITSCH and T. MANIATIS (1989): *Molecular cloning: a laboratory manual*, 2nd Edn, Cold Spring Harbour Laboratory Press, New York.
- STETTLER, R. F., H. D. BRADSHAW, P. E. HEILMAN and T. M. HINCKLEY (1996): *Biology of Populus: Implications for management and conservation*. National Research Council of Canada, Ottawa.
- STIRLING, B., G. NEWCOMBE, J. VREBALOV, I. BOSDET and H. D. BRADSHAW (2001): Suppressed recombination around the MXC3 locus, a major gene for resistance to poplar leaf rust. *Theor. Appl. Genet.* **103**: 1129–1137.
- STRONG, S. J., Y. OHTA, G. W. LITMAN and C. T. AMEMIYA (1997): Marked improvement of PAC and BAC cloning is achieved using electroelution of pulsed-field gel-separated partial digests of genomic DNA. *Nucleic Acids Res.* **25**: 3959–3961.
- TOMKINS, J. P., D. G. PETERSON, T. J. YANG, D. MAIN, E. F. ABLETT, R. J. HENRY, L. S. LEE, T. A. HOLTON, D. WATERS and R. A. WING (2001): Grape (*Vitis vinifera* L.) BAC library construction, preliminary STC analysis, and identification of clones associated with flavonoid and stilbene biosynthesis. *Amer. J. Enomol. Viticult.* **52**: 287–291.
- VILANOVA, S., C. ROMERO, D. ABERNATHY, A. G. ABBOTT, L. BURGOS, G. LLACER and M. L. BADENES (2003): Construction and application of a bacterial artificial chromosome (BAC) library of *Prunus armeniaca* L. for the identification of clones linked to the self-incompatibility locus. *Mol. Genet. Genomics* **269**: 685–691.
- VINATZER, B. A., H. B. ZHANG and S. SANSVINI (1998): Construction and characterization of a bacterial artificial chromosome library of apple. *Theor. Appl. Genet.* **97**: 1183–1190.
- WANG, W., W. ZHAI, M. LUO, G. JIANG, X. CHEN, X. LI, R. A. WING and L. ZHU (2001): Chromosome landing at the bacterial blight resistance gene Xa4 locus using a deep coverage rice BAC library. *Mol. Genet. Genomics* **265**: 118–125.
- XU, M. L., J. Q. SONG, Z. K. CHENG, J. M. JIANG and S. S. KORBAN (2001): A bacterial artificial chromosome (BAC) library of *Malus floribunda* 821 and contig construction for positional cloning of the apple scab resistance gene Vf. *Genome* **44**: 1104–1113.
- YIN, T. M., M. R. HUANG, M. X. WANG, L. H. ZHU, Z. B. ZENG and R. L. WU (2001): Preliminary interspecific genetic maps of the *Populus* genome constructed from RAPD markers. *Genome* **44**: 602–609.
- YIN, T. M., X. Y. ZHANG, M. R. HUANG, M. X. WANG, Q. ZHUGE, S. M. TU, L. H. ZHU and R. L. WU (2002): Molecular linkage maps of the *Populus* genome. *Genome* **45**: 541–555.
- YOKOYA, K., A. V. ROBERTS, J. MOTTLEY, R. LEWIS and P. E. BRANDHAM (2000): Nuclear DNA amounts in roses. *Ann. Bot.* **85**: 557–561.
- ZHU, H. and R. A. DEAN (1999): A novel method for increasing the transformation efficiency of *E. coli* – application for bacterial artificial chromosome library construction. *Nucleic Acids Res.* **27**: 910–912.