



Article High Temperature-Induced Spindle Destabilization Results in Aborted Pollen Production in *Populus*

Zhiqun Li ^{1,2,3}, Yifan Zhao ^{1,2,3}, Xuetong Cheng ^{1,2,3}, Bo Kong ^{1,2,3}, Yaru Sang ^{1,2,3}, Qing Zhou ^{1,2,3}, Jian Wu ^{1,2,3} and Pingdong Zhang ^{1,2,3,*}

- State Key Laboratory of Efficient Production of Forest Tree Resources, Beijing Forestry University, Beijing 100083, China
- ² Key Laboratory of Genetics and Breeding in Forest Trees and Ornamental Plants, Ministry of Education, Beijing Forestry University, Beijing 100083, China
- ³ School of Bioscience and Biotechnology, Beijing Forestry University, Beijing 100083, China
- * Correspondence: zhangpd@bjfu.edu.cn

Abstract: High temperature can induce the production of 2n gametes and aborted pollen during microsporogenesis in *Populus canescens*. However, the mechanism by which high temperature induces pollen abortion remains unknown. Here, pollen abortion was induced by exposing male flower buds of *P. canescens* to 38 and 41 °C; pollen morphology, meiotic abnormalities, defects of the meiotic microtubular cytoskeleton, and tapetum development were characterized. We found that the dominant meiotic stage, temperature, and duration of treatment significantly affected the percentage of high temperature-induced aborted pollen. Damaged spindle microtubules and depolymerized microtubular cytoskeletons were observed, which resulted in many lagging chromosomes at anaphase I and II as well as aneuploid male gametes and micronuclei, thereby generating aborted pollen grains. Tapetum disintegration was also delayed. The anther dehisced normally, and some viable pollen grains were released. However, no sufficient evidence showed that the delayed degradation of the tapetum was responsible for pollen abortion.

check for updates

Citation: Li, Z.; Zhao, Y.; Cheng, X.; Kong, B.; Sang, Y.; Zhou, Q.; Wu, J.; Zhang, P. High Temperature-Induced Spindle Destabilization Results in Aborted Pollen Production in *Populus*. *Forests* **2023**, *14*, 797. https:// doi.org/10.3390/f14040797

Academic Editor: Rita Lourenço Costa

Received: 25 February 2023 Revised: 4 April 2023 Accepted: 11 April 2023 Published: 13 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). **Keywords:** high temperature; aborted pollen; spindle; microtubular cytoskeleton; tapetum; *Populus canescens*

1. Introduction

In the context of global climate change, the key climatic parameters seem to be adversely affected are average surface temperatures, atmospheric CO_2 concentrations, rainfall regimes, and ultraviolet radiation. Concerning the temperature, over the past 100 years, global air temperatures have increased by 0.74 ± 0.18 °C, and an increasing rate of warming has taken place over the last 25 years. Overall, global climate change models predict a temperature increasing ranging from 1.8 °C to 4.0 °C over the next century. Obviously, such environmental changes would have significant effects on plant growth and development, species distribution, and diversity [1,2]. On a whole-plant level, elevated temperature induces multiple physiological disruptions, affecting both vegetative and reproductive processes. The sexual reproductive stage seems to be particularly vulnerable, resulting in a crop yield reduction being the most negative expected effect [3,4].

In *Populus*, owing to the sensitivity of pollen mother cells (PMCs) to environmental temperature during meiosis, high temperature is often used as a physical mutagenic agent to induce 2n pollen [5–7]. Zhang et al. [6] showed that the maximum frequency of 2n pollen that could be induced by high temperature exposure in *P. alba* L. was 39.6%. Tian et al. [7] demonstrated that the highest frequency of induced 2n pollen (56.3%) was achieved in *Populus canescens* (Aiton) Smith. Among the above studies, except for 2n pollen, a large number of aborted pollen grains were commonly observed. However, the mechanism by which aborted pollen is produced by high temperature remains unknown.

Abortion is a common reproductive phenomenon that is closely related to the amount of pollen, pollen viability, fertilization, and growth [8]. Previous studies have identified the following cytological factors as causes of abortion: (i) meiosis, (ii) tapetum development and disintegration, and (iii) cytoskeleton formation and disintegration [9–11]. First, meiosis is particularly susceptible to environmental changes. Many cytological events including cytomixis and abnormal chromosome behaviors occur during meiosis. Cytomixis events involving the exchange of chromosomes between meiocytes could contribute to the formation of unbalanced tetrads and polyads, which partly explains pollen abortion [12]. In addition, abnormal chromosome behaviors contribute to aborted pollen. Second, abortion is generally associated with tapetum development and disintegration. On the one hand, tapetum cells might secrete some components such as callose and sporopollenin precursors, which contribute to the dissolution of callose and the formation of exine, respectively [13]. The decrease in callose and sporopollenin might inhibit the development of microspores, leading to pollen abortion [14]. On the other hand, the abnormal enlargement and disintegration of the tapetum can result in a lack of nutrition for microspores and thus lead to abortion [10]. Finally, the cytoskeleton is responsible for mediating dynamic changes during meiosis and thus is closely associated with abortion. For example, aberrant cytoskeletal dynamics and an abnormal distribution of the cytoskeleton can ultimately lead to pollen abortion [11].

After flower buds are treated with high temperature in higher plants, macrospores or eggs form and mature in the ovule, which can only be observed through paraffin sectioning. Consequently, determining the effects of high temperature exposure on the fertility of female gametes is difficult. However, evaluating the effects of high temperature exposure on the fertility of pollen grains after they are released from anthers is less of a challenge by comparison. The objective of this study was to characterize the mechanism by which pollen abortion is induced via high temperature in *P. canescens*. Pollen abortion was induced by temperatures of 38 and 41 °C; pollen morphology, meiotic abnormalities, defects of the microtubular cytoskeleton during meiosis, and tapetum development were characterized, and expression analysis of *PtActin* gene was conducted. Our findings provide new insights into the mechanisms of heat stress-induced pollen abortion.

2. Materials and Methods

2.1. Plant Materials

Male floral branches ($2n = 2 \times = 38$) of *P. canescens* were collected from a natural population in Aletai, Xinjiang Uygur Autonomous Region, China. All sampled male branches were trimmed and cultured in tap water in a greenhouse (10–20 °C) located in Beijing Forestry University to promote floral development. No nutrients were added to the tap water.

2.2. High Temperature Exposure and Detection of Aborted Pollen

High temperature exposure was conducted according to the methods described by Tian et al. [7] Male flower buds were sampled at the following dominant meiotic stages: leptotence, zytotence, pachytence, diplotence, diakinesis, and metaphase I. The male flower buds were exposed to temperatures of 38 and 41 °C for 3 h and 6 h in a chamber, respectively. The air humidity of the chamber was 70%. Untreated male flower buds were used as the control group. After each treatment, all treated flower branches continued to be hydroponically cultured until anthers matured. Pollen samples were collected and stored in tubes with allochronic silica gel. To detect aborted pollen, some pollen grains were sampled, spread on a microscopic slide with a needle, and mounted in a drop of aceto-carmine solution (2%). Pollen grains were stained with acetocarmine solution, and unstained pollen grains were considered aborted pollen.

For scanning electron microscopy analysis, fixed pollen grains were rinsed with ethanol and air-dried. Samples were sputter-coated with gold using a HITACHI E-1010 ion sputter coater and observed under a HITACHIS-3400N microscope (Hitachi Limited, Tokyo, Japan), with an accelerating voltage of 5 kV.

For evaluation of induced pollen viability, a pollen germination test was conducted in vitro according to the methods reported by Zhao et al. [15] The medium for pollen germination had a pH value of 6.0. Pollen germination was conducted in a climate chamber at 26 °C. At 6 h after culture, the sampled pollen grains were fixed in Carnoy's fluid for 10 min after being centrifuged for 5 min at a speed of 1000 rpm. The germination rate of each sample was calculated using an eyepiece micrometre. A total of 300–600 pollen grains were assessed per sample. The germination rate was estimated by the number of germinated pollen grains and the total number of sampled pollen grains.

2.3. Meiotic Analysis of Pollen Mother Cells

As previously described by Tian et al. [7], when all treatments were finished, two to three flower buds were randomly sampled from the control group and the treatment groups every 2 h and fixed in Carnoy's fluid (ethanol:acetic acid, 3:1) until tetrads appeared. After 24 h, the fixed male buds were transferred to a 70% ethanol solution and stored in a refrigerator at 4 °C. For meiosis analysis, some anthers were dissected from the fixed buds using forceps and were crushed in a drop of aceto-carmine solution (2%) onto a microscope slide. Photomicrographs of developing microsporocytes (characterized by different meiotic stages) were placed under a fluorescence microscope (model BX51; Olympus, Tokyo, Japan) with a CCD camera (model DP70; Olympus, Tokyo, Japan). A total of 200–300 PMCs were counted per sample.

2.4. Tubulin Immunolocalization

The fresh anthers were randomly sampled from the control group and treatment groups and fixed in 4% paraformaldehyde (PEM buffer, pH 6.9) for 60 min. Next, α -tubulin immunolocalization analysis of PMCs during meiosis was performed as described previously [16]. Photomicrographs of developing microsporocytes (characterized by different meiotic stages) were collected with a Leica TCS-SP8 confocal laser scanning microscope.

2.5. Observation of Tapetum Development during Microspore Maturation

When PMCs were at the tetrad stage, two to three flower buds were randomly sampled from the control and treated groups every two days and fixed in Carnoy's fluid. After 24 h, the fixed male buds were transferred to the 70% ethanol solution and stored in a refrigerator at 4 °C. For observation of tapetum development, some anthers were dissected from the fixed buds using forceps and dehydrated in alcohol and xylene, embedded with paraffin, and sectioned for 8 μ m. The sections were stained with toluidine blue and photographed under an Olympus BX51 microscope(Olympus, Tokyo, Japan).

2.6. Statistical Analysis

We used a general linear model (GLM) to determine the effect of meiotic stage, temperature, and duration of treatment on the abortion rate of pollen. Data for the abortion rate of induced pollen were transformed (1/p) before analyses of variance to achieve homogeneous variances. All statistical analyses were conducted with SPSS software (SPSS for Windows, Version 13, SPSS, Chicago, IL, USA).

3. Results

3.1. The Effects of High Temperature on Aborted Pollen Production, Pollen Morphology and Pollen Viability

Male buds of *P. canescens* developed slowly and anthers became dry and brown after high temperature treatment. Some buds were dead after they were exposed to high temperatures, which precluded pollen collection.

High temperature-induced pollen was collected from all surviving male buds. A high proportion of aborted pollen was observed in addition to some induced 2n pollen

(Figure 1a). Spontaneously aborted pollen was rarely observed in the control group (Figure 1b). The percentage of aborted pollen in different treatments is shown in Table 1. The average percentage of high temperature-induced aborted pollen varied from 9.45 to 25.11%. GLM-univariate analysis of the percentages of high temperature-induced aborted pollen showed that dominant meiotic stage (F = 5.75, p < 0.001), temperature (F = 25.71, p < 0.001), and duration of treatment (F = 17.82, p < 0.001) significantly affected the percentages of high temperature-induced aborted pollen. Dominant meiotic stage \times temperature (F = 3.96, p = 0.004) and dominant meiotic stage \times duration (F = 5.48, p < 0.001) interactions also significantly affected the percentage of high temperature-induced aborted pollen. Temperature \times duration (F = 2.40, *p* = 0.128) and dominant meiotic stage \times temperature \times duration (F = 1.21, p = 0.319) interactions had no significant effect on the percentage of high temperature-induced aborted pollen. LSD multiple comparison tests showed that the differences in induced aborted pollen production were significantly higher at the diplotene stage than at the leptotene, pachytene, diakinesis, and metaphase I stages ($\alpha = 0.05$). The percentage of high temperature-induced aborted pollen was higher at 41 °C than at 38 °C. The percentage of high temperature-induced aborted pollen was higher for samples that were treated after 6 h than after 3 h.



Figure 1. Aborted pollen and 2n pollen induced via high temperature exposure and spontaneously aborted pollen in *P. canescens* (scale bar = $20.0 \mu m$). (a) Aborted pollen (arrow) and high temperature-induced 2n pollen (arrowhead) in *P. canescens*. (b) Spontaneously aborted pollen (arrow) in *P. canescens*.

Hours after Being Cultured	Dominant Meiotic Stage of PMCs	Temperature (°C)	Duration (h)	Percentage of Aborted Pollen (%)
52	Leptotene	38	3	9.45 ± 0.92
		38	6	17.15 ± 2.56
		41	3	11.29 ± 1.77
		41	6	17.92 ± 2.55
64	Zygotene	38	3	12.78 ± 5.06
		38	6	22.45 ± 2.21
		41	3	17.39 ± 5.65
		41	6	20.37 ± 0.83
76	Pachytene	38	3	15.72 ± 2.02
		38	6	15.35 ± 4.33
		41	3	18.61 ± 5.20
		41	6	16.16 ± 1.35

 Table 1. Aborted pollen production induced by high temperature in P. canescens.

Hours after Being Cultured	Dominant Meiotic Stage of PMCs	Temperature (°C)	Duration (h)	Percentage of Aborted Pollen (%)
88	Diplotene	38	3	19.90 ± 3.03
		38	6	19.33 ± 0.96
		41	3	20.97 ± 4.10
		41	6	20.93 ± 1.70
94	Diakinesis	38	3	14.17 ± 1.80
		38	6	13.83 ± 1.73
		41	3	16.95 ± 1.82
		41	6	20.62 ± 3.49
100	Metaphase I	38	3	10.82 ± 3.41
		38	6	13.76 ± 2.06
		41	3	25.11 ± 4.28
		41	6	24.13 ± 4.05
	Control			7.58 ± 1.32

Table 1. Cont.

Pollen morphology was studied by scanning electron microscopy. The pollen grains within the control group was uniform and spherical, with few corrugations and a granular surface (Figure 2a–c). There were no apertures on the surface. Few aborted pollen grains (arrows) were induced (Figure 2b,c) by 38 °C high temperature for 3 h or 6 h. The morphology (Figure 2d,e) and the ectexine structure (Figure 2f–i) of induced pollen were similar to the control group, suggesting that high temperature had no significant effects on the pollen morphology.



Figure 2. Scanning electron micrographs of pollen grains of *P. canescens* male flower buds exposed to high temperature (38 °C for 3 h or 6 h) or 25 °C (control) at the diplotene stage. (a) Morphology of pollen grains from the control group. (b) Ectexine structure of pollen grains from the control group.

(c) Details of the ectexine structure in pollen of the control group. (d) Morphology of pollen grains exposed to high temperature (38 °C for 3 h) (white arrows show aborted pollen grains). (e) Morphology of pollen grains exposed to high temperature (38 °C for 6 h) (white arrows show aborted pollen grains). (f) Ectexine structure of pollen exposed to high temperature (38 °C for 6 h). (g) Ectexine structure of pollen exposed to high temperature (38 °C for 6 h). (h) Details of the ectexine structure in pollen exposed to high temperature (38 °C for 6 h). (h) Details of the ectexine structure in pollen exposed to high temperature (38 °C for 3 h). (i) Details of the ectexine structure in pollen exposed to high temperature (38 °C for 6 h). (h) Details of the ectexine structure in pollen exposed to high temperature (38 °C for 6 h). (i) Details of the ectexine structure in pollen exposed to high temperature (38 °C for 6 h). (ii) Details of the ectexine structure in pollen exposed to high temperature (38 °C for 6 h). (b) f,g), and 5 µm (c,h,i).

For evaluation of induced pollen viability, fresh high temperature-induced pollen grains were supplied for germination testing on a medium containing 0.7% agar, 50 mg/L calcium chloride, 120 mg/L boric acid. Some germinated pollen grains were, respectively, observed for the control (Figure 3a), treatment at 38 °C for 3 h (Figure 3b) and treatment at 38 °C for 6 h (Figure 3c) groups. Pollen germination rates are shown in Figure 3d. After 6 h of culture, the average germination rate in the control group was 26.95%, which was slightly higher than that of induced pollen after treatments at 38 °C for 3 h (21.52%) and at 38 °C for 6 h (20.29%), indicating that the frequency of aborted pollen in the treatment groups were slightly higher than that in the control group. Among the treatments, the average germination rate of pollen subjected to 38 °C treatment for 3 h was slightly higher than that of pollen subjected to 38 °C treatment for 3 h was slightly higher than that of pollen subjected to 38 °C treatment for 3 h was slightly higher than that of pollen subjected to 38 °C treatment for 3 h was slightly higher than that of pollen subjected to 38 °C treatment for 3 h was slightly higher than that of pollen subjected to 38 °C treatment for 3 h was slightly higher than that of pollen subjected to 38 °C treatment for 6 h. However, the GLM-univariate analysis of germination rates revealed that high temperatures had no significant effect on induced pollen germination rates.



Figure 3. Effect of high temperature on pollen viability in *P. canescens.* (**a**) Germinated fresh pollen grains derived from the control group. (**b**) Germinated pollen grains derived from the treatment at 38 °C for 3 h. (**c**) Germinated pollen grains derived from the treatment at 38 °C for 6 h. (**d**) Pollen viability in flower buds exposed to 38 °C for 3 or 6 h in *P. canescens.* Control corresponds to 25 °C, that is, standard culture conditions. Bars = 50 μ m.

3.2. Induced Meiotic Abnormalities by High Temperature

In male flower buds that were exposed to 38 °C for 3 h or 6 h at the diploptene stage, all treated flower branches continued to be hydroponically cultured until pollen was released from the anthers. At the anaphase I, anaphase II, and tetrad stages, five flower buds per treatment were sampled to determine the effect of high temperature on the meiosis of PMCs. In the control group, the PMCs underwent normal meiosis, and a few lagging chromosomes in some PMCs were observed at anaphase I (Figure 4a) or anaphase II (Figure 4b). The number of lagging homologous chromosomes was $22.7 \pm 4.2\%$ at anaphase I. The number of lagging sister chromosomes varied from 1 to 10, and the percentage of PMCs with lagging sister chromosomes was $13.3 \pm 5.0\%$ at anaphase II (Table 2).



Figure 4. High temperature-induced lagging chromosomes and aberrant cytokinesis during the microsporogenesis of *P. canescens.* (a) Anaphase I in a control PMC. (b) Lagging chromosomes (arrow) at anaphase II in a control PMC. (c) Lagging chromosomes (arrow) in anaphase I derived from a PMC previously exposed to 38 °C for 3 h. (d) Lagging chromosomes (arrow) in anaphase II derived from a PMC previously exposed to 38 °C for 3 h. (e) Lagging chromosomes (arrow) in anaphase I derived from a PMC previously exposed to 38 °C for 6 h. (f) Lagging chromosomes (arrow) in anaphase I derived from a PMC previously exposed to 38 °C for 6 h. (f) Lagging chromosomes (arrow) in anaphase I derived from a PMC previously exposed to 38 °C for 6 h. (f) Lagging chromosomes (arrow) in anaphase II derived from a PMC previously exposed to 38 °C for 6 h. (g) Normal tetrad in a control PMC. (h) Polyad derived from a PMC previously exposed to 38 °C for 3 h. (i) Polyad with unbalanced cytokinesis derived from a PMC previously exposed to 38 °C for 6 h. Bars = 10 μ m.

Duration (h)	Percentage of PMCs with Lagging Chromosomes (%)		Number of Lagging Chromosomes	
	Anaphase I	Anaphase II	Anaphase I	Anaphase II
3	$40.67\pm3.06~^{\rm b}$	$33.33 \pm 8.00 \ ^{b}$	1–13	1–20
6	$55.33\pm4.16^{\text{ b}}$	48.67 ± 3.06 ^b	1–28	1–23
Control	$22.67\pm4.16~^{\rm a}$	13.33 ± 5.03 $^{\rm a}$	1–8	1–10

Table 2. Formation of lagging chromosomes induced by high temperature exposure in *P. canescens*.

Note: Lowercase letters indicate significance at the p < 0.05 level.

In some PMCs exposed to 38 °C, meiosis was abnormal, and a large number of lagging chromosomes were observed at anaphase I or II (Figure 4c–f). After 3 h of treatment at 38 °C, the number of lagging homologous chromosomes varied from 1 to 13, and the percentage of PMCs with lagging homologous chromosomes was $40.7 \pm 3.1\%$ at anaphase I. The number of lagging sister chromosomes varied from 1 to 20, and the percentage of PMCs with lagging homologous or sister chromosomes was significantly higher after treatment at 38 °C for 3 h compared with the control group. When PMCs were treated at 38 °C for 6 h, the number of lagging homologous chromosomes varied from 1 to 28, and the percentage of PMCs with lagging sister chromosomes with lagging homologous chromosomes was $55.3 \pm 4.2\%$ at anaphase I. The number of lagging sister chromosomes was $48.7 \pm 3.1\%$ at anaphase II (Table 2). The percentage of PMCs with lagging sister chromosomes was $48.7 \pm 3.1\%$ at anaphase II (Table 2). The percentage of PMCs with lagging sister chromosomes was $48.7 \pm 3.1\%$ at anaphase II (Table 2). The percentage of PMCs with lagging sister chromosomes was $48.7 \pm 3.1\%$ at anaphase II (Table 2). The percentage of PMCs with lagging homologous or sister chromosomes was $48.7 \pm 3.1\%$ at anaphase II (Table 2). The percentage of PMCs with lagging homologous or sister chromosomes was slightly higher after treatment at 38 °C for 6 h compared with PMCs treated at 38 °C for 3 h.

After two rounds of cell division, normal cytokinesis occurred in the control group, resulting in tetrad formation (Figure 4g). However, few micronuclei were observed in some of the treated PMCs (Figure 4h,i), and some polyads formed at the tetrad stage, suggesting that chromosome segregation errors occurred and led to the production of aneuploid gametes.

3.3. Delayed Tapetum Development Is Not Responsible for Induced Aborted Pollen Production

Defective tapetum development is often associated with the disrupted development of meiocytes and/or pollen and reduced/impaired fertility [17]. We examined the effect of high temperature on tapetum development via sectioning and toluidine blue staining during microspore maturation. At the tetrad stage in the control group, the tapetum was observed in the anther, which surrounded the developing PMCs (Figure 5a). Two days after the tetrad stage, the tapetum degenerated normally by programmed cell death (PCD) (Figure 5b). Four days after the tetrad stage, thinner tapetal layer cells were observed in the anther (Figure 5c). Six days after the tetrad stage, no tapetal layer cells were observed in the anther (Figure 5d). Eight days after the tetrad stage, the anther dehisced, and a large number of pollen grains with good fertility were released (Figure 5e).

At the tetrad stage in the group exposed to 38 °C for 3 h, the tapetum was observed in the anther, which surrounded the developing PMCs (Figure 5f). Two days after the tetrad stage, tapetum degenerated gradually by PCD (Figure 5g). Four days after the tetrad stage, thinner tapetal layer cells were observed in the anther (Figure 5h). Six days after the tetrad stage, few tapetal layer cells were observed in the anther (Figure 5i), suggesting that the tapetum of the treated anthers degenerated more slowly than the tapetum in the control group. Eight days after the tetrad stage, the anther matured, and more abortive pollen grains were released (Figure 5j). A similar pattern of tapetum degeneration was observed for PMCs exposed to 38 °C for 6 h (Figure 5k–n), except that many more abortive pollen grains were released after anthers matured (Figure 5o).



Figure 5. Tapetum development of anthers of *P. canescens* male flower buds exposed to high temperature (38 °C for 3 h or 6 h) or 25 °C (control). (**a**–**e**) Tapetum development of anthers from the control group. (**f**–**j**) Tapetum development of anthers exposed to 38 °C for 3 h. (**k**–**o**) Tapetum development of anthers exposed to 38 °C for 6 h. Scale bars = 5 μ m.

3.4. Induced Spindle Destabilization Results in Aborted Pollen Production

Because the meiotic microtubular cytoskeleton plays an important role in the segregation of homologous and sister chromosomes during meiosis, we examined the integrity and localization of microtubule structures of meiocytes exposed to 38 °C for 3 or 6 h using tubulin- α immunocytology. Under control conditions, the microtubular cytoskeleton was distributed regularly in microsporocytes (Figure 6a–h), which allowed for the proper segregation of chromosomes in daughter cells. After PMCs were treated at 38 °C for 3 h or 6 h at the diplotene, all meiotic stages exhibited the same microtubular distribution pattern (Figure 6i–x), similar to unstressed meiocytes (Figure 6a–h). However, the high temperature treatment resulted in some defects in the meiotic microtubular cytoskeleton. For example, the microtubules depolymerized and were not present in the cytoplasm. The extent of the treatment affected the degree of microtubule destabilization, and the plasma membrane even ruptured in some cells.

In the metaphase of the first meiotic division, a single bipolar spindle microtubule depolymerized in cells treated at 38 °C for 3 h (Figure 6j) and was damaged in cells treated at 38 °C for 6 h (Figure 6r). The damaged spindle microtubule resulted in the slow movement of chromosomes within the cytoplasm. This is consistent with the large number of lagging homologous chromosomes observed at anaphase I (Figure 4d,g). Similar to metaphase II, a double bipolar spindle microtubule at metaphase II depolymerized in cells treated at 38 °C for 3 h (Figure 6n) and was damaged in cells treated at 38 °C for 6 h (Figure 6v). Subsequently, several lagging sister chromosomes at anaphase II were observed (Figure 4e,h).

Under control conditions, telophase II male meiocytes typically generate a microtubule network that consists of six tetrahedrally arranged phragmoplast-like structures, which are localized between the four haploid nuclei (Figure 6h). However, telophase II male meiocytes treated at 38 °C for 3 h generated a new complex microtubule network that surrounded more than four nuclei instead of the six tetrahedrally arranged phragmoplast-like structure (Figure 6p). Microtubule bundles between two nuclei were much thinner in cells treated at 38 °C for 6 h (Figure 6x) than cells treated at 38 °C for 3 h and control cells.



Figure 6. Cont.



Figure 6. Defects in the meiotic microtubular cytoskeletons of PMCs exposed to 38 °C for 3 h or 6 h in *P. canescens.* (**a**–**h**) Meiotic microtubule distributions of PMCs from the control group. (**i**–**p**) Defects in the meiotic microtubular arrangement of PMCs after treatment at 38 °C for 3 h. (**q**–**x**) Defects in the meiotic microtubular arrangement of PMCs after treatment at 38 °C for 6 h. (**a**,**i**,**q**) Diakinesis. (**b**,**j**,**r**) Metaphase I. (**c**,**k**,**s**) Anaphase I. (**d**,**l**,**t**) Telophase I. (**e**,**m**,**u**) Prophase II. (**f**,**n**,**v**) Metaphase II. (**g**,**o**,**w**) Anaphase II. (**h**,**p**,**x**) Telophase II. Tubulin- α is shown in green, and DAPI is shown in blue. Scale bars = 10 µm.

4. Discussion

Heat stress (32 to 40 °C) will decline pollen viability. Many former studies have showed a decrease in the number of produced and released pollen grains [18,19], a reduction in pollen viability [20], as well as changes in both the pollen wall structure [18,20] and female fertility [21,22]. Pollination in flowering plants sometimes were affected by the high temperature. Koti et al. [18] documented that exposing to high day/night temperatures (38/30 °C) affected the pollen tube growth in soybean during flowering. When flowering *Brassica napus* was treated at 35 °C for 4 h, a reduction in the pollen germination rate was observed by Young et al. [22]. Cross et al. reported that the process of fertilization in flax

was affected by cyclical (daytime high 40 °C and night-time low 18 °C) heat stress 12 days after the initiation of flowering [23]. Subsequent events, such as embryogenesis and seed maturation, were also shown to be potentially sensitive to heat stress, leading to progenies with increased leaf area or delayed bud formation and dehardening [24–26].

In our study, high temperatures (38 and 41 °C) were shown to be capable of inducing the production of a large amount of aborted pollen. Suitable high temperature exposure affected the amount of aborted pollen in both a dosage- and meiotic stage-dependent manner. Flower buds had higher amounts of aborted pollen when exposed to higher temperatures. Heat exposure during microsporogenesis has been suggested to lead to pollen abortion in various species such as *Phaseolus vulgaris* [20], cotton [27], *Oryza sativa* [28], *Solanum lycopersicum* [29,30], and wheat [31,32]. However, in *P. canescens*, this response is only observed during a narrow window of development (limited to meiosis), which was consistent with the findings of in *Rosa* [8].

Meiosis is a specialized type of cell division that gives rise to daughter cells with reduced numbers of chromosomes. The reliable segregation of chromosomes is particularly important during meiosis, which is the specialized cell division that results in the formation of eggs and sperm. The egg and the sperm need to have one copy of each chromosome to give rise to a healthy embryo upon fertilization. If normal chromosome segregation is disrupted by an inhibitor of the spindle at anaphase I and II, chromosome segregation errors during meiotic division lead to the production of aneuploid gametes.

Given rapid changes in global climate, the effects of temperature on plant reproduction have received increased interest [3]. The process of meiosis of PMCs is sensitive to high temperature stress. Therefore, rising global temperatures threaten global crop production [33]. Previous studies have shown that heat stress results in numerous meiotic abnormalities, including chromosome stickiness, laggards, micronuclei, and spindle disorientation in the second meiotic division and aberrant cytokinesis, which suggests that meiosis in higher plants is sensitive to heat stress [34,35]. Here, male flower buds of *P. canescens* at the diplotene stage during the first cell division were exposed to 38 °C for 3 or 6 h. After high temperature treatment, chromosome segregation was delayed, and several lagging chromosomes were observed at anaphase I and II, resulting in the formation of aneuploid gametes and micronuclei. Some polyads were observed at the tetrad stage. Therefore, the aborted development of male meiocytes was observed prior to meiotic cytokinesis. This is consistent with observations of aborted pollen induced by 38 °C treatment in previous experiments [7].

The tapetum is located at the interface between gametophytic and sporophytic tissues and provides important elements for the development of male gametophyte, such as callose or sporopollenin precursors to help callose dissolution and exine formation, respectively [13]. The process of the tapetum's development can generally be divided into three stages: tapetum specification, tapetal cells' binucleation, and degeneration through PCD [13]. The fate of sexual cells and anther somatic cells is determined at floral stage 8/anther stage 4. However, both the tapetum and PMCs form at floral stage 9 and anther stage 5 in *Arabidopsis* [36], respectively. Defective tapetum development is often related to the disrupted development of meiocytes and/or pollen and reduced/impaired fertility [17,37,38]. For instance, the formation of the normal microspore cell wall and the fertility level can be affected by a lack of callose or a temporal shift in its activity. Jin et al. documented that a lack of callose production could lead to complete male sterility in Glycine max [14]. Early callose degradation was found to be sufficient for causing male sterility in *Nicotiana tabacum* [39].

Many previous studies have shown that heat stress can lead to male sterility in *Oryza* sativa L., Solanum melongena L., and Solanum lycopersicum Lam [40,41]. For example, in heat-susceptible barely plants, heat stress lasting four days or longer during the early phase of anther development caused the premature degradation of tapetum cells and PCD, leading to complete male sterility [42,43]. PCD is crucial for breaking down anther wall cells, such as in the tapetum and middle layer during pollen grain maturation and anther

dehiscence [44,45]. After male flower buds of *P. canescens* were treated at 38 °C for 3 or 6 h, the degradation of the tapetum was delayed. The anther dehisced normally, and some viable pollen grains were released. However, no sufficient evidence showed that the delayed degradation of the tapetum was responsible for the production of aborted pollen.

During microgenesis, both the movement of homologous or sister chromosomes and postmeiotic cytokinesis are closely related to dynamic changes in the organization of microtubules [46,47]. In diploid *Populus* species, the misorientation of spindle microtubules and the failure of postmeiotic cytokinesis have been attributed to the formation of unreduced gametes [6,16,35]. In Arabidopsis (*Arabidopsis thaliana*), callose and other cell plate components are deposited on the division planes by a network of microtubule (MT) arrays. The production of diploid male gametes in Arabidopsis can be induced by the cold-induced destabilization of post-meiotic radial microtubule arrays [48].

During cell division, chromosome segregation is considered to be driven by a spindle consisting of microtubules. Firstly, the microtubules capture and align the chromosomes at the spindle center. The chromosomes are segregated and moved to the poles of spindles during anaphase. The movement of chromosomes is pulled by the shortening of microtubule bundles which are linked to the chromosomes' kinetochores [49]. These kinetochore fibers (K-fibers) combine with a great number of microtubule-associated motor and nonmotor proteins to align and segregate the chromosomes [50].

A previous study showed that heat stress induced the depolymerization of the meiotic microtubular cytoskeleton, resulting in the failure of chromosome segregation in *Populus*. The microtubular cytoskeleton was able to repolymerize in some heat-treated cells after they were placed under normal conditions. However, aberrant cytokinesis occurred because of defects in the new radial microtubule systems, leading to the production of monads, dyads, triads, and polyads [51]. In our study, high temperature induced the depolymerization of meiotic microtubular organization and damaged the spindle microtubules, resulting in homologous chromosome and sister chromosome segregation errors at anaphase I and II. Some chromosomes were retained in the cytoplasm of microsporocytes because of the lack of shortening of the microtubule bundles. Therefore, high temperature-induced spindle destabilization was responsible for aborted pollen production.

5. Conclusions

This study strengthens knowledge of the effects of high temperatures on the development of PMCs of *Populus* during meiosis. The dominant meiotic stage, temperature, and duration of treatment significantly affected the percentage of high temperature-induced aborted pollen. Damaged spindle microtubules and depolymerized microtubular cytoskeletons were observed, which resulted in many lagging chromosomes at anaphase I and II, as well as aneuploid male gametes and micronuclei, forming aborted pollen grains. Tapetum disintegration was also delayed. The anther dehisced normally, and some viable pollen grains were released. However, no sufficient evidence showed that the delayed degradation of the tapetum was responsible for pollen abortion.

Author Contributions: Conceptualization, P.Z.; methodology, Z.L.; software, Y.Z.; resources, X.C.; data curation, Y.S.; writing—original draft preparation, Z.L. and B.K.; writing—review and editing, Q.Z. and J.W.; project administration, P.Z.; All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by the National Key R&D Program of China during the 14th Five-year Plan Period (2022YFD2200301-02).

Data Availability Statement: All the data that support the findings of this study are provided in the submission.

Acknowledgments: We thank Meiqin Liu and Junna Shi for their scanning electron microscopy analysis. We also thank Christopher A. at the University of North Carolina at Chapel Hill for his language editing. This research was supported by the National Key R&D Program of China during the 14th Five-year Plan Period (2022YFD2200301-02).

Conflicts of Interest: The authors declare no conflict of interest.

References

- 1. Minorsky, P.V. Global warming: Effects on plants. Plant Phys. 2002, 129, 1421–1422. [CrossRef] [PubMed]
- 2. Porter, J.R.; Semenov, M.A. Crop responses to climatic variation. Philos. Trans. R. Soc. B 2005, 360, 2021–2035. [CrossRef]
- Hedhly, A.; Hormaza, J.I.; Herrero, M. Global warming and sexual plant reproduction. *Trends Plant Sci.* 2009, 14, 30–36. [CrossRef]
 [PubMed]
- 4. Khan, A.H.; Min, L.; Ma, Y.Z.; Zeeshan, M.; Jin, S.X.; Zhang, X.L. High-temperature stress in crops: Male sterility, yield loss and potential remedy approaches. *Plant Biotechnol. J.* **2022**, *21*, 680–697. [CrossRef]
- 5. Kang, X.Y.; Zhu, Z.T.; Zhang, Z.Y. Suitable period of high temperature treatment for 2n pollen of *Populus tomentosa* × *P. bolleana*. *J. Beijing For. Univ.* **2000**, *22*, 1–4. [CrossRef]
- Zhang, L.; Wang, J.; Suo, Y.J.; Kang, X.Y. Pollen chromosome doubling under high temperature in *Populus alba. J. Nucl. Agric. Sci.* 2010, 24, 1158–1165. [CrossRef]
- Tian, M.D.; Zhang, Y.; Liu, Y.; Kang, X.Y.; Zhang, P.D. High temperature exposure did not affect induced 2n pollen viability in Populus. Plant Cell Environ. 2018, 41, 1383–1393. [CrossRef] [PubMed]
- 8. Pécrix, Y.; Rallo, G.; Folzer, H.; Cigna, M.; Gudin, S.; Le Bris, M. Polyploidization mechanisms: Temperature environment can induce diploid gamete formation in *Rosa* sp. *J. Exp. Bot.* **2011**, *62*, 3587–3597. [CrossRef] [PubMed]
- 9. Guan, J.Z.; Wang, J.J.; Cheng, Z.H.; Liu, Y.; Li, Z.Y. Cytomixis and meiotic abnormalities during microsporogenesis are responsible for male sterility and chromosome variations in *Houttuynia cordata. Genet. Mol. Res.* **2012**, *11*, 121–130. [CrossRef]
- Liu, L.W.; Huang, H.; Gong, Y.Q.; Chen, C.S.; Wang, L.Z. Cytological and ultra-structural study on microsporogenesis of cytoplasmic male sterility in *Raphanus sativus*. J. Integr. Plant Biol. 2009, 51, 850–857. [CrossRef]
- 11. Xu, C.G.; Liu, Z.T.; Zhang, L.P.; Zhao, C.P.; Yuan, S.H.; Zhang, F.T. Organizaiton of actin cytoskeleton during meiosis I in a wheat thermo-sensitive genic male sterile line. *Protoplasma* **2013**, *250*, 415–422. [CrossRef] [PubMed]
- Singhal, V.K.; Kumar, P. Cytomixis during microsporogenesis in the diploid and tetraploid cytotypes of Withania somnifera (L.) dunal, 1852 (solanaceae). Comp. Cytogenet. 2008, 2, 85–92. [CrossRef]
- 13. Scott, R.J.; Spielman, M.; Dickinson, H.G. Stamen structure and function. Plant Cell 2004, 16, 46–60. [CrossRef]
- 14. Jin, W.; Horner, H.T.; Palmer, R.G. Genetics and cytology of a new genic male-sterile soybean [*Glycine max* (L.) Merr.]. Sex. Plant Reprod. **1997**, 10, 13–21. [CrossRef]
- 15. Zhao, C.G.; Tian, M.D.; Li, Y.J.; Zhang, P.D. Slow-growing pollen-tube of colchicine-induced 2n pollen responsible for low triploid production rate in *Populus*. *Euphytica* **2017**, *213*, 94. [CrossRef]
- 16. Zhang, P.D.; Kang, X.Y. Occurrence and cytological mechanism of numerically unreduced pollen in diploid *Populus euphratica*. *Silvae Genet.* **2013**, *62*, 285–291. [CrossRef]
- 17. Chen, Z.S.; Liu, X.F.; Wang, D.H.; Chen, R.; Zhang, X.L.; Xu, Z.H.; Bai, S.N. Transcription factor OsTGA10 is a target of the MADS protein OsMADS8 and is required for tapetum development. *Plant Physiol.* **2018**, *176*, 819–835. [CrossRef]
- 18. Koti, S.; Reddy, K.R.; Reddy, V.R.; Kakani, V.G.; Zhao, D.L. Interactive effects of carbon dioxide, temperature, and ultraviolet-B radiation on soybean (*Glycine max* L.) flower and pollen morphology, pollen production, germination, and tube lengths. *J. Exp. Bot.* **2005**, *56*, 725–736. [CrossRef]
- Sato, S.; Kamiyama, M.; Iwata, N.; Furukawa, H.; Ikeda, H. Moderate increase of mean daily temperature adversely affects fruit set of *Lycopersicon esculentum* by disrupting specific physiological processes in male reproductive development. *Ann. Bot.* 2006, 97, 731–738. [CrossRef]
- 20. Porch, T.G.; Jahn, M. Effects of high temperature stress on microsporogenesis in heat-sensitive and heat-tolerant genotypes of *Phaseolus vulgaris. Plant Cell Environ.* **2001**, *24*, 723–731. [CrossRef]
- Peet, M.M.; Sato, S.; Gardner, R.G. Comparing heat stress effects on male-fertile and male-sterile tomatoes. *Plant Cell Environ*. 1998, 21, 225–231. [CrossRef]
- Young, L.W.; Wilen, R.W.; Bonham-Smith, P.C. High temperature stress of *Brassica napus* during flowering reduces micro- and megagametophyte fertility, induces fruit abortion, and disrupts seed production. *J. Exp. Bot.* 2004, 55, 485–495. [CrossRef] [PubMed]
- 23. Cross, R.H.; McKay, S.A.B.; McHughen, A.G.; Bonham-Smith, P.C. Heat-stress effects on reproduction and seed set in *Linum usitatissimum* L. (flax). *Plant Cell Environ.* 2003, 26, 1013–1020. [CrossRef]
- 24. Johnsen, Ø.; Dæhlen, O.G.; Østreng, G.; Skrøppa, T. Day length and temperature during seed production interactively affect adaptive performance of *Picea* abies progenies. *New Phytol.* **2005**, *168*, 589–596. [CrossRef]
- Johnsen, Ø.; Fossdal, C.G.; Nagy, N.; Mølmann, J.; Dæhlen, O.G.; Skrøppa, T. Climatic adaptation in *Picea abies* progenies is affected by the temperature during zygotic embryogenesis and seed maturation. *Plant Cell Environ.* 2005, 28, 1090–1102. [CrossRef]
- Lacey, E.P.; Herr, D. Parental effects in plant agolanceolata L. III. Measuring parental temperature effects in the field. *Evolution* 2000, 54, 1207–1217. [CrossRef]
- 27. Li, Y.L.; Li, Y.Y.; Su, Q.; Wu, Y.L.; Zhang, R.; Li, Y.W.; Ma, Y.Z.; Ma, H.H.; Guo, X.P.; Zhu, L.F. High temperature induces male sterility via MYB66-MYB4-Casein kinase Isignaling in cotton. *Plant Physiol.* **2022**, *189*, 2091–2109. [CrossRef]

- Peng, G.Q.; Liu, Z.L.; Zhuang, C.X.; Zhou, H. Environment-sensitive genic male sterility in rice and other plants. *Plant Cell Environ.* 2023, 46, 1120–1142. [CrossRef]
- 29. Iwahori, S. High temperature injuries in tomato. IV. development of normal flower buds and morphological abnormalities of flower buds treated with high temperature. *Hort. J.* **1965**, *34*, 33–41. [CrossRef]
- Sato, S.; Peet, M.M.; Thomas, J.F. Determining critical pre- and post-anthesis periods and physiological processes in *Lycopersicon* esculentum Mill. exposed to moderately elevated temperatures. J. Exp. Bot. 2002, 53, 1187–1195. [CrossRef]
- 31. Yang, X.T.; Ye, J.L.; Niu, F.Q.; Feng, Y.; Song, X.Y. Identification and verification of genes related to pollen development and male sterility induced by high temperature in the thermo-sensitive genic male sterile wheat line. *Planta* **2021**, *4*, 83. [CrossRef] [PubMed]
- 32. Bokshi, A.I.; Tan, D.; Thistlethwaite, R.J.; Trethowan, R.; Kunz, K. Impact of elevated CO2 and heat stress on wheat pollen viability and grain production. *Funct. Plant Biol.* **2021**, *48*, 503–514. [CrossRef]
- 33. Santiago, J.P.; Sharkey, T.D. Pollen development at high temperature and role of carbon and nitrogen metabolites. *Plant Cell Environ.* **2019**, *42*, 2759–2775. [CrossRef]
- Bomblies, K.; Higgins, J.D.; Yant, L. Meiosis evolves: Adaptation to external and internal environments. *New phytol.* 2015, 208, 306–323. [CrossRef]
- Wang, J.; Kang, X.Y. Distribution of microtubular cytoskeletons and organelle nucleoids during microsporogenesis in a 2n pollen producer of hybrid *Populus*. *Silvae Genet*. 2009, *58*, 220–226. [CrossRef]
- Sanders, P.M.; Bui, A.Q.; Weterings, K.; McIntire, K.N.; Hsu, Y.C.; Lee, P.Y.; Truong, M.T.; Beals, T.P.; Goldberg, R.B. Anther developmental defects in *Arabidopsis thaliana* male sterile mutants. *Sex. Plant Reprod.* 1999, 11, 297–322. [CrossRef]
- Cao, H.; Li, X.; Wang, Z.; Ding, M.; Sun, Y.; Dong, F.; Chen, F.; Liu, L.; Doughty, J.; Li, Y.; et al. Histone H2B monoubiquitination mediated by HISTONE MONOUBIQUITINATION1 and HISTONE MONOUBIQUITINATION2 is involved in anther development by regulating tapetum degradation-related genes in rice. *Plant Physiol.* 2015, *168*, 1389–1405. [CrossRef]
- Yi, J.; Moon, S.; Lee, Y.S.; Zhu, L.; Liang, W.; Zhang, D.; Jung, K.H.; An, G. Defective Tapetum Cell Death 1 (DTC1) regulates ROS levels by binding to metallothionein during tapetum degeneration. *Plant Physiol.* 2016, 170, 1611–1623. [CrossRef] [PubMed]
- Worrall, D.; Hird, D.L.; Hodge, R.; Paul, W.; Draper, J.; Scott, R. Premature dissolution of the microsporocyte callose wall causes male sterility in transgenic tobacco. *Plant Cell* 1992, 4, 759–771. [CrossRef]
- Li, B.; Chen, X.P.; Wu, Y.R.; Gu, A.X.; Zhang, J.J.; Luo, S.X.; Gao, X.R.; Zhao, J.J.; Pan, X.Q.; Shen, S.X. Gene characterization and molecular pathway analysis of reverse thermosensitive genic male sterility in eggplant (*Solanum melongena* L.). *Hortic. Res.* 2019, 6, 118. [CrossRef]
- Endo, M.; Tsuchiya, T.; Hamada, K.; Kawamura, S.; Yano, K.; Ohshima, M.; Higashitani, A.; Watanabe, M.; Kawagishi-Kobayashi, M. High temperatures cause male sterility in rice plants with transcriptional alterations during pollen development. *Plant Cell Physiol.* 2009, 50, 1911–1922. [CrossRef]
- Abiko, M.; Akibayashi, K.; Sakata, T.; Kimura, M.; Kihara, M.; Itoh, K.; Asamizu, E.; Sato, S.; Takahashi, H.; Higashitani, A. Hightemperature induction of male sterility during barley (*Hordeum vulgare* L.) anther development is mediated by transcriptional inhibition. *Plant Reprod.* 2005, 18, 91–100. [CrossRef]
- Oshino, T.; Abiko, M.; Saito, R.; Ichiishi, E.; Endo, M.; Kawagishi-Kobayashi, M.; Higashitani, A. Premature progression of anther early developmental programs accompanied by comprehensive alterations in transcription during high temperature injury in barley plants. *Mol. Genet. Genom.* 2007, 278, 31–42. [CrossRef] [PubMed]
- 44. Papini, A.; Mosti, S.; Brighigna, L. Programmed-cell-death events during tapetum development of angiosperms. *Protoplasma* **1999**, 207, 213–221. [CrossRef]
- Varnier, A.L.; Mazeyrat-Gourbeyre, F.; Sangwan, R.S.; Clément, C. Programmed cell death progressively models the development of anther sporophytic tissues from the tapetum and is triggered in pollen grains during maturation. *J. Struct. Biol.* 2005, 152, 118–128. [CrossRef]
- 46. Shamina, N.V.; Gordeeva, E.I.; Kovaleva, N.M.; Seriukova, E.G.; Dorogova, N.V. Formation and function of phragmoplast during successive cytokinesis stages in higher plant meiosis. *Cell Biol. Int.* **2007**, *31*, 626–635. [CrossRef] [PubMed]
- Brown, R.C.; Lemmon, B.E. Minispindles and cytoplasmic domains in microsporogenesis of Orchids. Protoplasma 1989, 148, 26–32.
 [CrossRef]
- 48. De Storme, N.; Copenhaver, G.P.; Geelen, D. Production of diploid male gametes in arabidopsis by cold-induced destabilization of postmeiotic radial microtubule arrays. *Plant Physiol.* **2012**, *160*, 1808–1826. [CrossRef]
- 49. Welburn, J.P.; Grishchuk, E.L.; Backer, C.B.; Wilson-Kubalek, E.M.; Yates, J.R.; Cheeseman, L.M. The human kinetochore Ska1 complex facilitates microtubule depolymerization-coupled motility. *Dev. Cell* **2009**, *16*, 374–385. [CrossRef]
- 50. Maiato, H.; DeLuca, J.; Salmon, E.D.; Earnshaw, W.C. The dynamic kinetochore-microtubule interface. J. Cell Sci. 2004, 117, 5461–5477. [CrossRef]
- 51. Wang, J.; Li, D.L.; Shang, F.N.; Kang, X.Y. High temperature-induced production of unreduced pollen and its cytological effects in *Populus. Sci. Rep.* 2017, *7*, 5281. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.