



BRAZILIAN JOURNAL
OF MEDICAL AND BIOLOGICAL RESEARCH

www.bjournal.com.br

ISSN 0100-879X
Volume 45 (2) 93-178 February 2012

BIOMEDICAL SCIENCES
AND
CLINICAL INVESTIGATION

Braz J Med Biol Res, February 2012, Volume 45(2) 118-124

doi: 10.1590/S0100-879X2012007500008

Identification of Albizia lebbeck seed coat chitin-binding vicilins (7S globulins) with high toxicity to the larvae of the bruchid *Callosobruchus maculatus*

A.J. Souza, A.T.S. Ferreira, J. Perales, D.G. Beghini, K.V.S. Fernandes, J. Xavier-Filho, T.M. Venancio and A.E.A. Oliveira

The Brazilian Journal of Medical and Biological Research is partially financed by



Ministério da Ciência e Tecnologia



Ministério da Educação



Institutional Sponsors



Explore High - Performance MS Orbitrap Technology In Proteomics & Metabolomics



analitica Thermo SCIENTIFIC
analiticaweb.com.br



All the contents of this journal, except where otherwise noted, is licensed under a [Creative Commons Attribution License](http://creativecommons.org/licenses/by-nc/4.0/)

Identification of *Albizia lebbbeck* seed coat chitin-binding vicilins (7S globulins) with high toxicity to the larvae of the bruchid *Callosobruchus maculatus*

A.J. Souza¹, A.T.S. Ferreira², J. Perales², D.G. Beghini², K.V.S. Fernandes¹,
J. Xavier-Filho¹, T.M. Venancio¹ and A.E.A. Oliveira¹

¹Laboratório de Química e Função de Proteínas e Peptídeos, Centro de Biociências e Biotecnologia, Universidade Estadual do Norte Fluminense, Campos dos Goytacazes, RJ, Brasil

²Laboratório de Toxinologia, Departamento de Fisiologia e Farmacodinâmica, Instituto Oswaldo Cruz, Fundação Oswaldo Cruz, Rio de Janeiro, RJ, Brasil

Abstract

Seed coat is a specialized maternal tissue that interfaces the embryo and the external environment during embryogenesis, dormancy and germination. In addition, it is the first defensive barrier against penetration by pathogens and herbivores. Here we show that *Albizia lebbbeck* seed coat dramatically compromises the oviposition, eclosion and development of the bruchid *Callosobruchus maculatus*. Dietary supplementation of bruchid larvae with *A. lebbbeck* seed coat flour causes severe weight loss and reduces survival. By means of protein purification, mass spectrometry and bioinformatic analyses, we show that chitin-binding vicilins are the main source of *A. lebbbeck* tegumental toxicity to *C. maculatus*. At concentrations as low as 0.1%, *A. lebbbeck* vicilins reduce larval mass from 8.1 ± 1.7 (mass of control larvae) to 1.8 ± 0.5 mg, which corresponds to a decrease of 78%. Seed coat toxicity constitutes an efficient defense mechanism, hindering insect predation and preventing embryo damage. We hypothesize that *A. lebbbeck* vicilins are good candidates for the genetic transformation of crop legumes to enhance resistance to bruchid predation.

Key words: Chitin-binding proteins; Seed coat; *Callosobruchus maculatus*; Insect bruchid

Introduction

During fruit development, seeds from several plants accumulate proteins, forming a critical nutritional source that is metabolized during germination and seedling growth (1). In eudicot seeds, 7S vicilins (globulins) are the main reserve proteins (1,2). In addition to their major role as energy source, alternative functions for vicilins have been proposed in sucrose binding (3), desiccation (4) and defense against insects and fungi (5-8). Several cotyledonary and tegumental vicilins can confer protection against *Callosobruchus maculatus* (9-12), the sugarcane borer *Diatraea saccharalis* (13) and *Tenebrio molitor* (8) larvae. In addition, vicilins isolated from various legume seeds (e.g., *Vigna angularis*, *Phaseolus vulgaris* and *Glycine max*) also impaired *C. maculatus* larval development (14). Furthermore, the mechanism of vicilin toxicity to some insects has been associated with their chitin-binding properties (5,6,12-14). Chitin is an extracellular amino-sugar polymer

of N-acetyl- β -glucosamine that is an essential structural component of fungal cell walls, exoskeletons and peritrophic membranes (PM; in arthropods) (15). PM is a non-cellular membrane consisting of chitin, proteins, glycoproteins, and proteoglycan conjugates (16). This structure has important roles in the recycling of digestive enzymes and in the protection of the epithelium against food particle abrasion and microorganism infection (16). *C. maculatus* predation seriously compromises the replanting and commercialization potential of *Vigna unguiculata* (cowpea) seeds, a key legume crop in the semi-arid tropics. In view of its critical agronomic relevance, we investigated the toxicity of *Albizia lebbbeck* seed coat proteins to *C. maculatus*. *A. lebbbeck* is a widespread species of the *Albizia* genus and recent studies have suggested its pharmacological potential as a source for anti-tumor, anti-fungal and anti-inflammatory molecules (17,18).

Correspondence: A.E.A. Oliveira, Av. Alberto Lamego, 2000, P5 Sala 224, 28013-602 Campos dos Goytacazes, RJ, Brasil.
E-mail: elenir@uenf.br

Received May 7, 2011. Accepted January 16, 2012. Available online January 27, 2011. Published February 17, 2012.

Material and Methods

Seeds and insects

A. lebbeck seeds were collected at Universidade Estadual do Norte Fluminense, Campos dos Goytacazes, RJ, Brazil. Commercial cowpea (*V. unguiculata* cv. fradinho) seeds were purchased from local markets at Campos dos Goytacazes, RJ, Brazil. *C. maculatus* (Coleoptera: Bruchidae) adults were maintained in an in-house colony and reared on cowpea seeds (cv. fradinho) at 28°C, 60-80% relative humidity, and a 12-h light:12-h dark photoperiod.

Seed infestation

Sixty *A. lebbeck* seeds were placed in glass flasks and exposed to ten 3-day-old *C. maculatus* females for 24 h. The experiment was conducted in a BOD incubator chamber at 28°C and 70% relative humidity. Females were removed and larval development was monitored using a stereoscopic microscope coupled to a digital CCD video camera, from oviposition to complete seed coat perforation or larval death. Control experiments using cowpea as host seeds were performed in parallel under the same conditions.

Extraction and purification of seed coat proteins

Seed coats were separated from cotyledons and ground with a pestle and mortar. Proteins were extracted (1:10 flour to buffer ratio) with 100 mM phosphate buffer with 500 mM NaCl, pH 7.6, for 3 h at 4°C and centrifuged at 10,000 *g* for 30 min. The supernatant was treated with 90% saturation with ammonium sulfate for 24 h at 4°C and centrifuged at 10,000 *g* for 30 min at 4°C. The precipitated fraction was dialyzed against water for 24 h at 4°C using 7-kDa cut-off membranes and freeze-dried.

The freeze-dried material was diluted (500 mg/46 mL 0.1 M sodium acetate buffer, pH 5.0) and fractionated by CM-Sephacryl ion exchange chromatography. The sample was applied to a 2.0 x 21.0 cm column equilibrated with 0.1 M sodium acetate buffer, pH 5.0. The non-retained fraction (CI) was eluted and adsorbed proteins were sequentially desorbed using 0.25 and 0.5 M NaCl solutions (CII and CIII fractions, respectively).

Gel filtration chromatography was used to separate a 50 mg/mL (50 mM potassium phosphate buffer, pH 7.8) sample of CI on a Sephacryl S-100 column (2.0 x 70 cm). The fractions (SI, SII, and SIII) were eluted using the sample dissolution buffer. The SI fraction was used for protein identification and characterization and toxicity to insects.

Protein identification and characterization

The protein fraction obtained by Sephacryl S-100 gel filtration chromatography (SI fraction) was analyzed by 15% SDS-polyacrylamide gel electrophoresis (SDS-PAGE) (19), conducted at 15 mA. Gels were stained with 0.05% Coomassie blue and destained in 10% acetic acid.

In-gel tryptic digestion of SI fraction proteins was per-

formed as described (20). For mass spectrometry analysis the peptides were co-crystallized with 0.3 µL 10 mg/mL α-cyano-4-hydroxycinnamic acid solution in 0.1% (w/v) trifluoroacetic acid, 50% (v/v) acetonitrile directly onto a MALDI target plate. Raw data for protein identification were obtained with a 4700 Proteomics Analyzer (Applied Biosystems, USA). Both mass spectrometry (MS) and MS/MS data were acquired in the positive and reflectron modes using a neodymium-doped yttrium aluminum garnet (Nd:YAG) laser with a 200-Hz repetition rate. Typically, 1600 shots were accumulated for spectra in the MS mode while 3000 shots were accumulated for spectra in the MS/MS mode. Up to 10 of the most intense ion signals with a signal-to-noise ratio above 20 were selected as precursors for MS/MS acquisition. Peptides from trypsin autolysis peaks and matrix ion signals were discarded. External calibration in the MS mode was performed using a mixture of four peptides: des-Arg1-bradykinin (*m/z* = 904.47), angiotensin I (*m/z* = 1296.69), Glu1-fibrinopeptide B (*m/z* = 1570.68), and ACTH (18-39) (*m/z* = 2465.20). MS/MS spectra were externally calibrated using fragments of known ion masses observed in the MS/MS spectrum of angiotensin I. MS/MS results were searched against the NCBI nr databases using the Mascot software (www.matrixscience.com). The peptide sequences were analyzed using the Scaffold 3™ software to determine peptide identification probability. Bit scores were computed by aligning the MS peptides using BLAST. Search parameters were as follows: two tryptic missed cleavages allowed, non-fixed modifications of methionine, tryptophan, histidine (oxidation), cysteine (carbamidomethylation and propionamide). The pps and ppw files were generated from the raw (or native) MS data according to the following parameters using the Data Explorer Software (Applied Biosystems, USA). For MS1: mass range: 900-4000; peak density: 15 peaks per 200 Da; signal-to-noise ratio: 20; minimum area: 100, and maximum peaks per spot: 60. For MS2: mass range: 60 and precursor 20; peak density: 55 peaks per 200 Da; signal-to-noise ratio: 2; minimum area: 10, and maximum peaks per precursor: 200.

Bioinformatic analyses

We used a computational strategy based on BLAST searches (command line version) (21) to identify the proteins (or close homologs from other species) from which the MS peptides were derived. Since the sequenced peptides are short, we used non-default BLAST parameters (word size 2, no filters or compositional based statistics and PAM30 as score matrix). The full-length amino acid sequences of the aligned hits were submitted to a second BLAST search, which allowed us to obtain a more comprehensive list of homologs, not biased by the short alignments from MS peptides. Sequences aligning with an E-value <0.001, 40% similarity and 50% query and hit coverage were recovered, clustered using blastclust (21), and aligned using MUSCLE (22). BLAST results were parsed using BioPerl (23) and custom Perl scripts. Taxonomic information was obtained from the Taxonomy database (NCBI) (24) using custom scripts.

Feeding trials

To test the effects of the seed coat flours and chromatographic fractions on larval development, we used an artificial seed system (9). Samples were added to a cowpea-based meal at concentrations of 0.1, 0.25, 0.5, and 1.0%. Artificial seeds were exposed to 3-day-old fertilized females for 24 h (28°C, 70% relative humidity). Females were removed and laid eggs were left at a concentration of 3 eggs per seed. After 20 days, infested seeds were opened and larvae were counted and weighed. Control experiments were performed under the same conditions using *V. unguiculata* (cv. fradinho) cotyledons. Experiments consisted of 3 seeds per assay and were run in triplicate (a total of 9 seeds and 27 eggs per tested dose). Statistical significance was assessed using the Student *t*-test (*P* value <0.05) (25).

ELISA

Vicilin levels were quantified by enzyme-linked immunosorbent assay (ELISA) (26), using an anti-*V. unguiculata* (cv. EPACE-10) vicilin antibody produced in rabbits (primary antibody) diluted 1:1000 and a peroxidase-conjugated anti-rabbit IgG antibody (secondary antibody) diluted 1:2000. Cotyledonary EPACE-10 vicilin samples (5 to 0.0024 µg/100 µL) were used as reference. Peroxidase activity was developed using ortho-phenylenediamine.

Chitin-binding assay

Chitin from lobster shells was obtained from Sigma-Aldrich (USA) and treated with a previously established protocol (27). The finely ground chitin was used to prepare an affinity column, subsequently used to purify chitin-binding proteins from the CI (162 mg/2 mL) and SI (18 mg/mL) fractions (dissolved in 80 mM sodium acetate buffer, pH 5.0). The non-retained fraction was eluted with the dilution buffer and adsorbed proteins were desorbed using 0.1 M HCl. Both fractions were dialyzed against water for 24 h at 4°C and freeze-dried.

Results

Toxicity of *A. lebeck* seed coat to *C. maculatus*

By comparing the survival and development of *C. maculatus* on *A. lebeck* and *V. unguiculata* cv. fradinho (host seed) we found that oviposition, larval eclosion and adult emergence were drastically compromised in the former. The reduction of oviposition was striking (96%), whereas seed coat penetration, larval eclosion and adult emergency were not observed before 30 days after oviposition. The amount of adults that emerged from the *V. unguiculata* seeds after this time was about 86% (data not shown).

Additional experiments using flour prepared from *A. lebeck* seed coats showed a clear dose-dependent toxicity to *C. maculatus* larvae, with strong weight loss and mortality by 20 days (Figure 1). At concentrations of 0.25, 0.5, and 1%, we observed 40, 83.8, and 87.1% larval weight loss,

respectively. In addition, after 20 days of incubation, only 14.1% of the larvae survived on a diet with 1% of *A. lebeck* seed coat flour (*P* < 0.05, *t*-test).

Purification and characterization of seed coat proteins and toxicity to *C. maculatus*

A. lebeck seed coat proteins were initially precipitated using 0-90% ammonium sulfate and submitted to ion exchange chromatography on a CM-Sephrose column, resulting in a non-retained fraction (CI) and two adsorbed fractions that were desorbed with 0.25 and 0.5 M NaCl (CII and CIII, respectively). Higher NaCl concentrations (0.75 and 1.0 M NaCl) did not desorb additional protein fractions (Figure 2). The three chromatographic fractions




Figure 1. Toxicity of *Albizia lebeck* seed coat flour to *Callosobruchus maculatus* larval development and survival at 20 days after oviposition. The seed coat protein was incorporated into artificial seeds at different relative concentrations (0.1, 0.25, 0.5, and 1.0%). Experiments were carried out in triplicate and the data shown are the means \pm SD of these results. **P* < 0.05 compared to control (*t*-test).




Figure 2. CM-Sephrose ion exchange chromatography (2.0 x 21.0 cm) of *Albizia lebeck* seed coat proteins previously precipitated with ammonium sulfate (0-90% saturation) and dissolved in sodium acetate buffer, pH 5.0. The column was equilibrated and chromatography was developed with 0.1 M sodium acetate buffer, pH 5.0. The non-retained fraction was eluted with 0.1 M sodium acetate buffer, pH 5.0 (CI fraction) and adsorbed proteins were desorbed with 0.25 and 0.5 M NaCl (CII and CIII fractions, respectively).

were incorporated into the larval diet and CI was the one with highest toxicity, with 67 and 100% larval mortality at concentrations of 0.1 and 0.25%, respectively (Figure 3A). In addition, we noticed that larval death can happen either before or after hatching (Figure 3B, Panels a-d). When the larval development on natural seeds of *A. lebeck* and *V. unguiculata* (*C. maculatus*-host seeds) was monitored, during the initial 6 days after oviposition no morphological differences in larval development were observed between the two seeds (Figure 3B, Panels e-j). Egg content was clear and larvae were virtually formed by the fifth day, when the larvae started to hatch and perforate the seed coat (Figure 3B, Panels e,f). Although they stayed alive for up to 16 days when feeding on *A. lebeck* (Figure 3B, Panel g), *C. maculatus* larvae were unable to reach the cotyledonary tissue (Figure 3B, Panel h). On the other hand, *C. maculatus* larvae were able to reach and intensively feed on the *V. unguiculata* cotyledons by the 6th day after oviposition (Figure 3B, Panels i,j).

Although CI administration resulted in a clear reduction of larval counts, the body weight of surviving individuals was apparently unaffected (Figure 3C). CII and CIII were also toxic at 1%, both showing a 67% mortality rate (Figure 3A) and CIII also reduced larval body weight by 77.2% (Figure 3C). Therefore, these results suggest that *A. lebeck* seed coat has physicochemically distinct sets of toxic proteins. Due to its higher toxic potential against *C. maculatus*, CI was elected for additional characterization experiments.

CI was submitted to Sephacryl S-100 gel filtration chromatography, resulting in three fractions (SI, SII, and SIII; Figure 4A). Due to its high abundance, we chose SI for further experiments. When the SI protein profile was examined by SDS-PAGE, three major bands with molecular masses between 50 and 40 kDa (Figure 4B) were demonstrable. These three major bands were cut out, subjected to trypsin digestion and MS analysis. The results showed that two peptides originated from bands 1 and 8 were derived from band 3 (Table 1). The sequences obtained were analyzed by BLAST in order to identify the protein. The peptides were identified as components of vicilin from *V. unguiculata* (ID GenBank emb|CAP19902.1) and *V. radiata* 8S globulin (ID GenBank ABG02260). Col-

lectively, the peptides account for approximately 24.7% of the *V. radiata* 8S protein and 25.9% of *V. unguiculata* vicilin sequences. No peptide was obtained from band 2 (45-kDa




Figure 3. Toxicity of flour and of the CI fraction prepared from *Albizia lebeck* seed coat to *Callosobruchus maculatus* larvae. **A** and **B**, Larval mortality. **B**, a-d: larval mortality in artificial seeds containing the CI fraction (arrows). a: the larva died inside the egg; b: perforation of artificial seed; c and d: dead larvae found inside the artificial seed. e-j: development of larvae on natural *A. lebeck* and *Vigna unguiculata* seeds. e: larva inside the egg on the 5th day after oviposition; f: inferior surface of the egg on the 5th day after oviposition; g: larva inside the egg on the 16th day after oviposition; h: *A. lebeck* seed coat partially perforated on the 16th day after oviposition; i: hole across the host seed coat from *V. unguiculata* on the 6th day; j: hole on the *V. unguiculata* cotyledon surface on the 6th day after oviposition. The bar inside **Panel a** indicates magnification for **Panels a-d**. The bar inside **Panel e** indicates magnification for **Panels e-g**. The bar inside **Panels h/i** indicates magnification for **Panels h-j**. **C**, Larval mass at 20 days after oviposition. The seed coat fraction was incorporated into artificial seeds at different concentrations (0.1, 0.25, 0.5, and 1.0%). Experiments were done in triplicate and the data shown are the means \pm SD of these results. * $P < 0.05$ compared to control (*t*-test).

region). The peptide sequences REQQQQQEESWE VQRY and RLHEITPEKNPQLRD obtained from protein band 3 were analyzed using the Scaffold 3™ software and the results showed a peptide identification probability of 95% with *V. unguiculata* vicilin (ID GenBank 160332746).

We used a computational strategy based on BLAST (21) to identify homologs in the nr database (see Methods for details). Although the peptides sequenced from each band were totally different from each other, the hit lists from the two bands had an extremely high overlap (more than 96% of the BLAST hits were shared by the bands). Therefore, our results strongly suggest that the two bands are isoforms of the same gene (or extremely close paralogs), and hence we merged the results. The combined list is dominated by vicilins and vicilin-like proteins (e.g., gis 160332746 and 145207915), which were extracted from the nr database, clustered using blastclust and aligned [data shown at (<http://dl.dropbox.com/u/7886304/supplements/BJMBR-Souza2012/BJMBR-Souza-MSA.pdf>)]. We extracted taxonomic information for all the hits and found that all the detected homologs were from the Tracheophyta clade (vascular plants), encompassing homologs in the basal vascular plant *Selaginella moellendorffii* (Lycopodiophyta), the fern *Matteuccia struthiopteris*, conifers, various monocots (e.g., *Oryza sativa* and *Zea mays*) and eudicots (e.g., *Vitis vinifera*, *Phaseolus vulgaris* and *Glycine max*) [data shown at (<http://dl.dropbox.com/u/7886304/supplements/BJMBR-Souza2012/BJMBR-Souza-MSA.pdf>)].

com/u/7886304/supplements/BJMBR-Souza2012/BJMBR-Souza-MSA.pdf]. Although homologs could be detected in such diverse major groups, our results indicate a proliferation of vicilins in eudicots, which is coherent with their prominent nutritional roles in such these seeds.

The SI fraction showed strong toxicity to *C. maculatus* larval development. At a concentration of 0.1%, larval mass was reduced from 8.1 ± 1.7 (mass of control larvae) to 1.8 ± 0.5 mg, which corresponds to a decrease of 78% (data not shown).




Figure 4. A, Sephacryl S-100 gel filtration chromatography of the CI fraction of *Albizia lebeck* seed coat previously obtained by CM-Sepharose chromatography resulting in three fractions (SI, SII, and SIII). The column was equilibrated and developed with 50 mM potassium phosphate buffer, pH 7.8. B, SDS-polyacrylamide gel electrophoresis of the SI fraction (106 to 116 mL).

Table 1. Mass spectrometry analysis of the peptides from the fraction of *Albizia lebeck* seed coat obtained by Sephacryl S-100 gel filtration chromatography (SI fraction).

| Spot No. | Peptide sequence | Protein accession number and Blast bit score* | |
|----------|----------------------|---|---|
| | | <i>Vigna radiata</i> 8S globulin ABG02260 | <i>Vigna unguiculata</i> vicilin emb CAP19902.1 |
| Band 1 | KQIQNLENYRV | 36.3 | 40.1 |
| | RIPAGTTFFLVNPNNDNLRI | 63.0 | 54.5 |
| Band 3 | RWFHTLFRN | 36.3 | 40.1 |
| | KIPAGTTFFLVNPDNENLRI | 71.0 | 61.7 |
| | RAILTLVNPDGRD | 41.4 | 43.9 |
| | KQIQNLENYRV | 36.3 | 40.1 |
| | RGQNNPFYFDSRW | 45.6 | 52.8 |
| | RLHEITPEKNPQLRD | 42.6 | 52.8 |
| | REQQQQQEESWEVQRY | 53.2 | 62.2 |
| | KLAIPVNNPHRF | 40.1 | 40.1 |

*Bit scores were computed by aligning the MS peptides against the *Vigna radiata* 8S globulin protein ABG02260.1 (454 amino acids) and *Vigna unguiculata* vicilin protein (433 amino acids) using BLAST. Collectively, the peptides encompass approximately 24.7% of the *V. radiata* 8S protein and 25.9% of *V. unguiculata* vicilin. We used a 90% threshold for query coverage and 35 bit score (BLAST) to map the peptides on the vicilin and vicilin-like proteins.




Figure 5. Chitin-binding affinity of *Albizia lebeck* seed coat SI protein fraction and quantification of vicilin-like proteins in the non-retained and adsorbed chitin fractions (desorbed with 0.1 M HCl).

The vicilin concentration in the SI fraction (1.4 µg/mg) was ~5.6-fold higher than the one found in the CI fraction that was 0.25 µg/mg (data not shown). In addition, proteins from the SI fraction also showed strong chitin-binding activity and, after desorption with 0.1 M HCl, were recognized by anti-vicilin antibodies (Figure 5).

Discussion

Before reaching the cotyledons, bruchid larvae must pass through the seed coat, which may pose not only a physical, but also a chemical toxic barrier to counter the penetration process (10,28). For instance, *Acanthoscelides obtectus* larvae were unable to penetrate *P. lunatus* seed coats (29) and only 50% of them penetrated *P. vulgaris* teguments (30). The seed coat of *Vicia faba* prevented the penetration of about 55% of the *Callosobruchus chinensis* and *C. maculatus* larvae (31,32).

In the present study, we demonstrated that *C. maculatus* oviposition, hatching and adult development were drastically reduced when the bruchids were grown on *A. lebeck* seeds.

References

- Shewry PR. Plant storage proteins. *Biol Rev Camb Philos Soc* 1995; 70: 375-426.
- Muntz K, Belozersky MA, Dunaevsky YE, Schlereth A, Tiedemann J. Stored proteinases and the initiation of storage protein mobilization in seeds during germination and seedling growth. *J Exp Bot* 2001; 52: 1741-1752.
- Braun H, Czihal A, Shutov AD, Baumlein H. A vicilin-like seed protein of cycads: similarity to sucrose-binding proteins. *Plant Mol Biol* 1996; 31: 35-44.
- Baumlein H, Braun H, Kakhovskaya IA, Shutov AD. Seed storage proteins of spermatophytes share a common ancestor with desiccation proteins of fungi. *J Mol Evol* 1995; 41: 1070-1075.
- Gomes VM, Okorokov LA, Sales MP, Fernandes KV, Rose TL, Xavier FJ. Vicilin storage proteins inhibit yeast growth and glucose stimulated acidification of the medium by cells. *Folia Microbiol* 1997; 42: 224.
- Sales MP, Pimenta PP, Paes NS, Grossi-de-Sa MF, Xavier-Filho J. Vicilins (7S storage globulins) of cowpea (*Vigna unguiculata*) seeds bind to chitinous structures of the midgut of *Callosobruchus maculatus* (Coleoptera: Bruchidae) larvae. *Braz J Med Biol Res* 2001; 34: 27-34.
- Uchoa AF, DaMatta RA, Retamal CA, Albuquerque-Cunha JM, Souza SM, Samuels RI, et al. Presence of the storage seed protein vicilin in internal organs of larval *Callosobruchus maculatus* (Coleoptera: Bruchidae). *J Insect Physiol* 2006; 52: 169-178.
- Paes EV, Uchoa AF, Pinto MST, Silva CP, Fernandes KVS, Oliveira AEA, et al. Binding of *Vigna unguiculata* vicilins to the peritrophic membrane of *Tenebrio molitor* affects larval development. *Entomol Exp Appl* 2008; 129: 11-17.
- Macedo MLR, Andrade LBD, Moraes RA, Xavier-Filho J. Vicilin variants and the resistance of Cowpea (*Vigna unguiculata*) seeds to the Cowpea Weevil (*Callosobruchus maculatus*). *Comp Biochem Physiol, C - Pharmacol Toxicol Endocrinol* 1993; 105: 89-94.

A. lebeck seed coat flour was shown to be toxic to the larvae, causing considerable lethality and weight loss of 20-day-old larvae. After several purification steps, we obtained a highly toxic vicilin-rich sample that was subsequently shown to have strong chitin-binding activity, with consequent toxicity of certain vicilins to chitin-containing structures in the insect midgut epithelium and fungal cell walls (5,8,12-14,33). It has been hypothesized that particular plant chitin-binding proteins can adhere to insect PMs, interfering with their normal functions and affecting nutrient absorption (12). In addition, chitin-binding vicilins from several plants have been demonstrated to be toxic to several species of insects and fungi (5,8,13,14,34). However, it has also been reported that intact vicilin and vicilin-derived peptides are absorbed by the insect's intestinal tract and incorporated into internal organs and eggs (7,35).

Distinct seeds are naturally predated by diverse bruchids, e.g., *V. unguiculata* by *C. maculatus* and *Phaseolus* by *Zabrotes subfasciatus*. This relative specificity and dietary preferences are molded by an evolutionary arm race between the insect and plant species (36). Taken together, our results and previous ones suggest that vicilin toxicity might be a major player in such evolutionary phenomenon and could thus be manipulated in plant engineering strategies aiming to enhance seed resistance to predators. However, the mechanism of toxicity of vicilins and their respective detoxification strategies by insects are yet to be described.

Acknowledgments

Research supported by FAPERJ, CNPq, and Universidade Estadual do Norte Fluminense Darcy Ribeiro.

10. Oliveira AEA, Machado OLT, Gomes VM, Neto JX, Pereira AC, Vieira JGH, et al. Jack bean seed coat contains a protein with complete sequence homology to bovine insulin. *Protein Peptide Lett* 2011; 6: 15-21.
11. Moraes RA, Sales MP, Pinto MS, Silva LB, Oliveira AE, Machado OL, et al. Lima bean (*Phaseolus lunatus*) seed coat phaseolin is detrimental to the cowpea weevil (*Callosobruchus maculatus*). *Braz J Med Biol Res* 2000; 33: 191-198.
12. Uchoa AF, de Miranda MR, de Souza AJ, Gomes VM, Fernandes KV, Lemos FJ, et al. Toxicity of hydrolyzed vicilins toward *Callosobruchus maculatus* and phytopathogenic fungi. *J Agric Food Chem* 2009; 57: 8056-8061.
13. Mota AC, DaMatta RA, Lima FM, Silva CP, Xavier-Filho J. Cowpea (*Vigna unguiculata*) vicilins bind to the peritrophic membrane of larval sugarcane stalk borer (*Diatraea saccharalis*). *J Insect Physiol* 2003; 49: 873-880.
14. Yunes ANA, de Andrade MT, Sales MP, Morais RA, Fernandes KVS, Gomes VM, et al. Legume seed vicilins (7S storage proteins) interfere with the development of the cowpea weevil (*Callosobruchus maculatus* (F)). *J Sci Food Agr* 1998; 76: 111-116.
15. Merzendorfer H, Zimoch L. Chitin metabolism in insects: structure, function and regulation of chitin synthases and chitinases. *J Exp Biol* 2003; 206: 4393-4412.
16. Terra WR. The origin and functions of the insect peritrophic membrane and peritrophic gel. *Arch Insect Biochem Physiol* 2001; 47: 47-61.
17. Lam SK, Ng TB. First report of an anti-tumor, anti-fungal, anti-yeast and anti-bacterial hemolysin from *Albizia lebbbeck* seeds. *Phytomedicine* 2011; 18: 601-608.
18. Babu NP, Pandikumar P, Ignacimuthu S. Anti-inflammatory activity of *Albizia lebbbeck* Benth., an ethnomedicinal plant, in acute and chronic animal models of inflammation. *J Ethnopharmacol* 2009; 125: 356-360.
19. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970; 227: 680-685.
20. Shevchenko A, Wilm M, Vorm O, Mann M. Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal Chem* 1996; 68: 850-858.
21. Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997; 25: 3389-3402.
22. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 2004; 32: 1792-1797.
23. Stajich JE, Block D, Boulez K, Brenner SE, Chervitz SA, Dagdigian C, et al. The Bioperl toolkit: Perl modules for the life sciences. *Genome Res* 2002; 12: 1611-1618.
24. Sayers EW, Barrett T, Benson DA, Bolton E, Bryant SH, Canese K, et al. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 2011; 39: D38-D51.
25. Bridge PD, Sawilowsky SS. Increasing physicians' awareness of the impact of statistics on research outcomes: comparative power of the t-test and Wilcoxon Rank-Sum test in small samples applied research. *J Clin Epidemiol* 1999; 52: 229-235.
26. Engvall E, Perlmann P. Enzyme-linked immunosorbent assay (ELISA). Quantitative assay of immunoglobulin G. *Immunochemistry* 1971; 8: 871-874.
27. Hackman RH, Goldberg M. New substrates for use with Chitinases. *Anal Biochem* 1964; 8: 397-401.
28. Silva LB, Sales MP, Oliveira AE, Machado OL, Fernandes KV, Xavier-Filho J. The seed coat of *Phaseolus vulgaris* interferes with the development of the cowpea weevil [*Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae)]. *An Acad Bras Cienc* 2004; 76: 57-65.
29. Thierry D. Hardness of some fabaceous seed coats in relation to larval penetration by *Acanthoscelides obtectus* (Say) (Coleoptera, Bruchidae). *J Stored Prod Res* 1984; 20: 177-181.
30. Thierry D, Jarry M, Pouzat J. To penetrate or not to penetrate - a behavioral choice by bean beetle 1st instar larvae in response to *Phaseolus vulgaris* seed surface quality. *J Chem Ecol* 1994; 20: 1867-1875.
31. Boughdad A, Gillon Y, Cagnepain C: effect of the ripe seed husks of vicia faba on the larval development of *Callosobruchus maculatus*. *Entomol Exp Appl* 1986; 42: 219-223.
32. Desroches P, Elshazly E, Mandon N, Duc G, Huignard J. Development of *Callosobruchus chinensis* (L) and *C. maculatus* (F) (Coleoptera, Bruchidae) in seeds of vicia faba L differing in their tannin, vicine and convicine contents. *J Stored Prod Res* 1995; 31: 83-89.
33. Sales MP, Gomes VM, Fernandes KV, Xavier-Filho J. Chitin-binding proteins from cowpea (*Vigna unguiculata*) seeds. *Braz J Med Biol Res* 1996; 29: 319-326.
34. Moura FT, Oliveira AS, Macedo LL, Vianna AL, Andrade LB, Martins-Miranda AS, et al. Effects of a chitin-binding vicilin from *Enterolobium contortisiliquum* seeds on bean bruchid pests (*Callosobruchus maculatus* and *Zabrotes subfasciatus*) and phytopathogenic fungi (*Fusarium solani* and *Colletrichum lindemuntianum*). *J Agric Food Chem* 2007; 55: 260-266.
35. Souza SM, Uchoa AF, Silva JR, Samuels RI, Oliveira AE, Oliveira EM, et al. The fate of vicilins, 7S storage globulins, in larvae and adult *Callosobruchus maculatus* (Coleoptera: Chrysomelidae: Bruchinae). *J Insect Physiol* 2010; 56: 1130-1138.
36. Vermeij GJ. *Evolution and escalation: an ecological history of life*. Princeton: Princeton University Press; 1987.